

Salmonella Serotypes - Non-human Isolates, October - December 2003

| Serotype | Animal | | | | | | | | Meat/bone meal | Environmental | Food | Spice | Not specified | Poultry | | | | Total |
|-----------------------------------|----------|-----------|----------|----------|----------|----------|----------|----------|----------------|---------------|-----------|----------|---------------|----------|-----------|---------------|---------------------------------|-------|
| | Avian | Bovine | Canine | Equine | Feline | Ovine | Porcine | Sealion | | | | | | Neckflap | Feed | Environmental | Miscellaneous including product | |
| Amsterdam | | | | | | | | | | 1 | | | | | | | 1 | |
| Anatum | | | | | | | | 5 | | | | | | | | | 5 | |
| Anatum 15+ | | | | | | | | | | | | | | | 1 | | 1 | |
| Brandenburg | | 4 | | | | | | 2 | 3 | 8 | | | | 4 | 3 | | 24 | |
| Cerro | | | | | | | 2 | | | | | | | | | | 2 | |
| Derby | | | | | | | 2 | | | | | | | | | | 2 | |
| Enteritidis phage type 9a | | | 1 | | | | | | | | | | | | | | 1 | |
| Give 15+ | | | | | | | | | | | | | | 2 | | | 2 | |
| Hadar | | | | | | | | | | | | | 1 | | | | 1 | |
| Havana | | | | | | | | 1 | 1 | | | | | 1 | 1 | | 4 | |
| Heidelberg | | 1 | | | | | | | | | | | | | | | 1 | |
| Hindmarsh | | 1 | | | | 2 | | | | 1 | | 2 | | | | | 6 | |
| Infantis | | | | | | | | | 1 | 5 | | | 1 | 3 | 1 | 1 | 12 | |
| Kentucky | | | | | | | | | | | | | | 1 | | | 1 | |
| London 15+ | | | | | | | | | | | | | | 1 | | | 1 | |
| Mbandaka | | | | | | | | | | | | | | 1 | | | 1 | |
| Montevideo | | | | | | | | | | 2 | | | | | | | 2 | |
| Newport | | | | | | | 5 | | | | | | | | | | 5 | |
| Orion 15+ | | | | | | | | | | 1 | | | | 2 | | | 3 | |
| Putten | | | | | | | | | | | | | | 1 | | | 1 | |
| Rubislaw | | | | | | | | | | | | 1 | | 1 | | 1 | 3 | |
| Senftenberg | | | | | | | | | | | | | | | 1 | | 1 | |
| Tennessee | | | | | | | | | 1 | | | 1 | | 2 | | | 4 | |
| Typhimurium phage type 1 | | 1 | | | | | | | | | | | | | | 1 | 2 | |
| 8 | | 2 | | | | | | | | | | 1 | | | | | 3 | |
| 9 | | 1 | | | | | | | | | | | 1 | | | | 2 | |
| 12a | | | | | | | | | | | | | | | 1 | 1 | 2 | |
| 23 | | 1 | | | | | | | | | | | | | | | 1 | |
| 42 | | | | | | | | | 17 | | | | | | | | 17 | |
| 74 | | | | | | | | | | | | | | 1 | | | 1 | |
| 101 | | 6 | | | | | | | | | | 1 | | | | | 7 | |
| 135 | 1 | 3 | | | | | | | | | | | 1 | | | | 5 | |
| 135 variant | | | | | | | | | | | | | 1 | | | 1 | 2 | |
| 155 | | | | | | 1 | | | | | | | | | | | 1 | |
| 156 | | 6 | 2 | | | | | | | | | | | | | | 8 | |
| 160 | | 1 | | 2 | 2 | | 1 | | 4 | 1 | | | | 1 | | 1 | 13 | |
| 193 | | | | | | | | | | | | | | 1 | | | 1 | |
| RDNC | 1 | | | | | 1 | | | | | | | | | | | 2 | |
| Untypable | | 2 | | | | | | | | | | | | 1 | | 2 | 5 | |
| Weltevreden 15+ | | | | | | | | | | 1 | | | | | | | 1 | |
| Group B 4,5,12 :-: - (non motile) | | | | | | | | | | | | 1 | | | | | 1 | |
| Group C 6,7 :-: - (non motile) | | | | | | | | | | | | | | | | 1 | 1 | |
| Group C 8, 20 :-: y :- | | | | | | | | | | 1 | | | | | | | 1 | |
| Group E 3,19 :-: - (non motile) | | | | | | | | | | | | | | | 1 | | 1 | |
| Group F 11 : r :- | | | | | | | | | | | | 1 | | | | | 1 | |
| TOTAL | 2 | 29 | 3 | 2 | 2 | 4 | 1 | 9 | 8 | 27 | 21 | 0 | 8 | 4 | 24 | 9 | 162 | |

S. Typhimurium 72
 Other serotypes 90
 Poultry isolates 46
 Animal isolates 52