

Salmonella Serotypes - Non-human Isolates, January-March 2004

| Serotype | Animal | | | | | | | | | | Meat/bone meal | Environmental | Food | Not specified | Poultry | | | | Sp Chicken * | Total |
|--------------------------------|----------|----------|----------|----------|----------|----------|-----------|----------|----------|----------|----------------|---------------|----------|---------------|----------|-----------|---------------|---------------------------------|--------------|------------|
| | Avian | Bovine | Canine | Cervine | Equine | Feline | Ovine | Porcine | Reptile | Sea Lion | | | | | Neckflap | Feed | Environmental | Miscellaneous including product | | |
| Adelaide | | | | | | | | | 2 | | | | | | | | | | | 2 |
| Agona | | | | | | | | | | | | | | | | 1 | 3 | | | 4 |
| Anatum | | | | | | | | | | | 3 | 1 | | | | 1 | | 5 | | 10 |
| Brandenburg | | | | 1 | | | | | | | 1 | | 2 | | | 1 | 3 | 1 | | 9 |
| Derby | | | | | | | | | | | | | | | | 17 | 1 | | | 18 |
| Enteritidis phage type 9a | | | | | | | 1 | | | | | | | | | | | 1 | | 2 |
| Havana | | | | | | | | | | | 2 | | | | | | 4 | | | 6 |
| Hindmarsh | | | | | | | 29 | | | | | | 1 | | | | | | | 30 |
| Infantis | | | 1 | | | | | | | | 1 | | | | 1 | | 2 | 1 | | 6 |
| Johannesburg | | | | | | | | | | | | 1 | | | | | | | | 1 |
| Kentucky | | | | | | | | | | | | | | | | | 1 | | | 1 |
| London | | | | | | | 1 | | | | | | | | | | | | | 1 |
| London 15+ | | | | | | | | | | | | | | | | 1 | | | | 1 |
| Mbandaka | | | | | | | | | | | | | 3 | | | | 1 | | | 4 |
| Montevideo | | | | | | | | | | | 1 | | | | | | | | | 1 |
| Muenchen | | | | | | | | | 4 | | | | | | | | | | | 4 |
| Orion 15+ | | | | | | | | | | | | | | | | | 1 | | | 1 |
| Paratyphi B var Java | | | | | | | | | | | | 2 | | | | | | | | 2 |
| Saintpaul | | 1 | 1 | | | | | | 1 | | | | | | | | | | | 3 |
| Senftenberg | | | | | | | | | | | | | | | | 3 | 1 | | | 4 |
| Tennessee | | | 1 | | | | | | | | 5 | | | | | | 6 | | | 12 |
| Thompson | 2 | | | | | | | | | | | | | | | | 1 | | | 3 |
| Typhimurium | | | | | | | | | | | | | | | | | | 2 | | 2 |
| Typhimurium phage type 1 | | | | | 1 | | 1 | | | | | | | | | | | | | 2 |
| 8 | | 1 | | | | | | | | | | | | | | | | | | 1 |
| 12a | | | | | | | | | 1 | | | | | | | | | | | 1 |
| 41 | | | | | | | | | | | 2 | | | | | | | | | 2 |
| 42 | | 1 | | | | | | | | | | | | | | | | 1 | | 2 |
| 101 | | | | | | | | | | | | | | | 1 | | 47 | 14 | | 62 |
| 101 variant | | | | | | | | | | | | | | | | | 2 | | | 2 |
| 135 | 3 | | | | | 1 | | 1 | | | | | 2 | | | | | | | 7 |
| 156 | | | | | 1 | | | | | | | | | | | 1 | 1 | | | 3 |
| 160 | 1 | 1 | 1 | | | 1 | | | | | | | | 8 | | 2 | 6 | 1 | | 21 |
| Untypable | | | | | | | | | | | | 1 | | | | | | | | 1 |
| Urbana | | | | | | | | | | | | 27 | | | | | | | | 27 |
| Weltevreden | | | | | | | | | | | | 1 | | | | | | | | 1 |
| Group B 4,12 :- : 1,2 | | | | | | | | | | | | | | | | 1 | | | | 1 |
| Group B 4,5,12 :- : - | | | | | | | | | | | | | | | | | | | 1 | 1 |
| Group C 6,7 :- : 1,5 | | | | | | | | | | | | | | | | | | 1 | | 1 |
| Group N 30 :- : - (non-motile) | | | | | | | | | | | | 12 | | | | | | | | 12 |
| TOTAL | 6 | 4 | 4 | 1 | 2 | 2 | 32 | 1 | 7 | 1 | 15 | 45 | 8 | 8 | 2 | 28 | 80 | 27 | 1 | 274 |

*This isolate was from a NZFSA special project

S. Typhimurium 106
 Other serotypes 167
 Poultry isolates 137
 Animal isolates 60