

Meningococcal Disease Report:
January 2009

Prepared for the Ministry of Health

By

Liza Lopez
Diana Martin

February 2009

Client Report
FW09012

Meningococcal Disease Report:

January 2009

Diana Martin
Principal Scientist
Programme and Project Leader, Communicable Disease

DISCLAIMER

This report or document (“the Report”) is given by the Institute of Environmental Science and Research Limited (“ESR”) solely for the benefit of the Ministry of Health, Public Health Service Providers and other Third Party Beneficiaries as defined in the Contract between ESR and the Ministry of Health, and is strictly subject to the conditions laid out in that contract.

Neither ESR nor any of its employees makes any warranty, express or implied, or assumes any legal liability or responsibility for use of the Report or its contents by any other person or organisation.

ACKNOWLEDGEMENTS

This report could not have been generated without the continuing support of staff in Public Health Services, clinical laboratories, medical practices and hospitals throughout New Zealand. All have a role in improving surveillance. The authors wish to especially thank Heather Davies for meningococcal specialist laboratory testing; Donald Peterkin and Carol Kliem for assistance with surveillance, and Dinusha Fernando for statistical advice.

TABLE OF CONTENTS

1. INTRODUCTION.....	4
2. METHODS	4
2.1. Surveillance Methods	4
2.2. Laboratory Methods.....	5
3. RESULTS.....	6
REFERENCES	17

LIST OF FIGURES AND TABLES

Figure 1: Meningococcal disease cases by month, 2005/2006 (average), 2007/2008 (average) and January 2009.....	6
Table 1. Geographic distribution by District Health Board of total and confirmed cases of meningococcal disease and proportion of confirmed to total cases, January 2005/2006 (average), 2007/2008 (average) and 2009.....	7
Table 2. Age distribution of total and confirmed cases of meningococcal disease and proportion of confirmed to total cases, January 2005/2006 (average), 2007/2008 (average) and 2009.....	8
Table 3. Meningococcal disease cases less than two years of age, January 2005/2006 (average), 2007/2008 (average) and 2009.....	8
Table 4. Meningococcal disease cases by District Health Board and age group, January 2005/2006 (average), 2007/2008 (average) and 2009.....	9
Table 5. Numbers and age-standardised incidence rates by ethnicity for cases of meningococcal disease and proportion of confirmed to total cases, January 2005/2006 (average), 2007/2008 (average) and 2009.....	10
Table 6. Meningococcal disease cases by strain type, January 2005/2006 (average), 2007/2008 (average) and 2009.....	10
Table 7. Meningococcal disease cases by strain type, February-January: 2007, 2008 and 2009.....	11
Table 8. Epidemic strain meningococcal disease by region, February-January: 2007, 2008 and 2009.....	11
Table 9. Epidemic strain meningococcal disease by age group, February-January: 2007, 2008 and 2009.....	11
Table 10. Epidemic strain meningococcal disease by ethnicity, February-January: 2007, 2008 and 2009.....	12
Table 11. Meningococcal disease cases by District Health Board and strain type, January 2005/2006 (average), 2007/2008 (average) and 2009.....	13
Table 12. Meningococcal disease cases by age group and strain type, January 2005/2006 (average), 2007/2008 (average) and 2009.....	14
Table 13. Meningococcal disease cases by basis for diagnosis, January 2005/2006 (average), 2007/2008 (average) and 2009.....	15
Table 14. Case-fatality rates for meningococcal disease cases by District Health Board, age, ethnicity, and group, January 2005/2006 (average), 2007/2008 (average) and 2009.....	16

1. INTRODUCTION

New Zealand's epidemic of serogroup B meningococcal disease began in mid-1991. In mid-2002 vaccine trials aimed at measuring immunogenicity and the safety of a 'tailor made' vaccine (MeNZB™) targeting New Zealand's epidemic 'strain' were started. In July 2004, provisional licensure was granted for the progressive rollout of MeNZB™ from 6 months to 19 years of age inclusive. The programme commenced in Counties Manukau DHB on 19 July 2004, and was progressively rolled out nationwide during 2005. Administration of the vaccine continues for those requiring completion of their doses.

This report aims to provide timely data on the epidemiology of meningococcal disease against which the effectiveness of the vaccine campaign can be assessed by comparing trends for 2009 against 2005/2006 average and 2007/2008 average for the same period. The initial report (July 2004) compared trends for January to June. Subsequent reports contain monthly updates.

2. METHODS

2.1. Surveillance Methods

Surveillance of meningococcal disease in New Zealand is based on a combination of notification and laboratory data. Meningococcal disease is notifiable to Medical Officers of Health under the Health Act 1956. Data on each case are recorded onto a computerised database (EpiSurv), which is installed in all Public Health Services (PHSs). These data are sent to ESR each week and collated on behalf of the Ministry of Health. Both patient specimens and meningococci or meningococcal DNA obtained from cases of disease are referred to the Meningococcus Reference Laboratory at ESR for confirmation of disease and for characterisation of the invading strain. Additionally, results from any diagnostic testing undertaken in clinical laboratories are actively sought for all notified cases. This information combined with group, PorB type (serotype) and PorA type (subtype), determined at ESR on isolates and identity of meningococcal DNA from patient specimens, are entered into a laboratory database and merged with the EpiSurv database to provide a more complete picture of the disease epidemiology.

Notification and laboratory data are based on information recorded as at 16 February 2009. Any changes made to either database after this date will not be reflected in this report. Excel 2003 was used to analyse the combined dataset. Rates expressed in this report are cumulative (as opposed to annualised). Hence in the initial report, rates were six-monthly, but in subsequent reports (updated monthly), they will be seven-monthly, eight-monthly etc. Therefore direct rate comparisons between the updated reports should not be made. Rates were calculated using 2006 population census data as the denominator for 2005/2006, 2007/2008 and 2009. The 2005/2006 and 2007/2008 results are averaged, rounded to one decimal place. Ethnicity was determined using the prioritised approach. Note: This report uses the 2006 census data which has impacted on ethnicity rates.

Earliest date (earliest of report date, onset date, hospitalisation date, date of death and laboratory dates) was used in this report to determine date of case.

Note: The numbers for 2008 are provisional and subject to change. The occurrence of late notifications and denotifications mean that the total for the year may change.

Case Definition

The case definition in the Ministry of Health's Communicable Disease Control Manual[1] is 'meningococcal disease presents as meningitis or meningococcal Septicaemia. The disease presents as an acute fever, nausea, vomiting, and headache, that may progress rapidly to shock and death. Petechial rash is seen in about 50 percent'. Cases with a clinically compatible illness are classified as confirmed or probable as follows:

Confirmed case: A clinically compatible illness with at least one of the following:

- isolation of *Neisseria meningitidis* from an otherwise sterile body site (blood, CSF, aspirate or skin biopsy); or
- a positive nucleic acid test (NAT) using polymerase chain reaction (PCR) on CSF, blood, serum, or aspirate; or
- detection of Gram-negative intracellular diplococci in CSF, aspirate or skin biopsy; or
- positive meningococcal antigen test on CSF.

Probable case:

- a clinically compatible illness and isolation of *N. meningitidis* from the throat; or
- a clinically compatible illness.

2.2. Laboratory Methods

All meningococci isolated and patient samples (blood and CSF) or DNA recovered from notified cases of meningococcal disease are referred from diagnostic laboratories in New Zealand to the Meningococcus Reference Laboratory (MRL) at ESR. The MRL then undertakes confirmation of the presence of a meningococcus or meningococcal DNA and characterisation to determine the strain type. This includes identification of the capsule group, PorB and PorA types.

Strain typing: The capsular group is identified either by the slide agglutination technique using commercial antisera specific for serogroups A, B, C, X, Y, Z, W135 and 29E or by PCR testing[2]. The PorB outer membrane protein (OMP) type, and the PorA OMP type are determined on isolates using the whole cell ELISA method of serotyping[3] or by PCR followed by sequence analysis[4]. For whole cell analysis the following Monoclonal antibodies (RIVM, The Netherlands, NIBSC, England) are used for detecting PorB antigens 1, 2a, 2b, 4, 14 and 15; and for PorA: P1.1, P1.2, P1.4, P1.5, P1.6, P1.7, P1.9, P1.10, P1.12, P1.13, P1.14, P1.15 and P1.16. Note the sequencing of the *porB* gene is not routinely undertaken.

The strain type is defined as, for example for the epidemic strain, B:4:P1.7-2,4 where B is the group, 4 is the PorB type and P1.7-2,4 defines the PorA type. Note the PorA type has two epitopes known as variable regions, which are identified. The 7-2 indicates there is a specific deletion in the VR1 epitope, otherwise defined as P1.7. Monoclonal antibodies do not recognise epitopes with deletions in the variable regions and hence would define this strain as B:4:P1.4.

Multi-locus sequence typing[5] is used to further characterise isolates of interest and when needed restriction fragment length polymorphism (RFLP) analysis of isolates[6] is carried out for the purpose of defining clusters.


B other are isolates that are group B but have PorA types that are distinguished from the epidemic type by serotyping/genotyping.

3. RESULTS

Meningococcal disease figures for January 2009 are eight notified cases with no deaths reported. The average figures for the same time period in 2007/2008 are nine notified cases and one death, the average for 2005/2006 being 15.5 notified cases and one death.

Meningococcal disease figures for the first month of 2009, and the average for the same time period in 2007/2008 and in 2005/2006 are presented below in graph and table format. As rates have been calculated even when the numerators are small, care should be taken when interpreting, most noticeably with the cumulative rate for some DHBs and the case fatality rates.

Figure 1: Meningococcal disease cases by month, 2005/2006 (average), 2007/2008 (average) and January 2009



Note: There is only one value for January 2009.

Table 1. Geographic distribution by District Health Board of total and confirmed cases of meningococcal disease and proportion of confirmed to total cases, January 2005/2006 (average), 2007/2008 (average) and 2009

District Health Board	Jan 2005/2006 average				Jan 2007/2008 average				Jan 2009			
	No. Total ¹	Rate ²	No. Conf ¹	% ³	No. Total ¹	Rate ²	No. Conf ¹	% ³	No. Total	Rate ²	No. Conf	% ³
Northland	0.5	0.3	0.5	100	0.5	0.3	0.5	100	0	0.0	0	-
Waitemata	3.5	0.7	1.5	42.9	1.0	0.2	1.0	100	0	0.0	0	-
Auckland	0.0	0.0	0.0	-	0.5	0.1	0.5	100	0	0.0	0	-
Counties Manukau	1.0	0.2	1.0	100	0.0	0.0	0.0	-	2	0.5	2	100
Waikato	0.5	0.1	0.5	100	0.0	0.0	0.0	-	0	0.0	0	-
Lakes	0.0	0.0	0.0	-	0.0	0.0	0.0	-	0	0.0	0	-
Bay of Plenty	0.5	0.3	0.5	100	0.0	0.0	0.0	-	1	0.5	1	100
Tairāwhiti	0.5	1.1	0.5	100	0.5	1.1	0.5	100	0	0.0	0	-
Taranaki	1.0	1.0	1.0	100	0.5	0.5	0.5	100	0	0.0	0	-
Hawke's Bay	1.0	0.7	1.0	100	2.0	1.3	1.5	75.0	1	0.7	1	100
Whanganui	0.0	0.0	0.0	-	0.0	0.0	0.0	-	0	0.0	0	-
MidCentral	1.0	0.6	1.0	100	0.0	0.0	0.0	-	0	0.0	0	-
Hutt	0.5	0.4	0.5	100	0.5	0.4	0.5	100	0	0.0	0	-
Capital and Coast	1.0	0.4	0.5	50.0	0.5	0.2	0.5	100	2	0.8	2	100
Wairarapa	0.5	1.3	0.0	0.0	0.0	0.0	0.0	-	0	0.0	0	-
Nelson												
Marlborough	1.0	0.8	1.0	100	0.5	0.4	0.5	100	0	0.0	0	-
West Coast	0.0	0.0	0.0	-	0.0	0.0	0.0	-	0	0.0	0	-
Canterbury	2.0	0.4	2.0	100	1.5	0.3	1.0	66.7	1	0.2	0	0.0
South Canterbury	0.0	0.0	0.0	-	0.0	0.0	0.0	-	0	0.0	0	-
Otago	0.5	0.3	0.5	100	0.0	0.0	0.0	-	1	0.6	1	100
Southland	0.5	0.5	0.5	100	1.0	0.9	1.0	100	0	0.0	0	-
New Zealand total	15.5	0.4	12.5	80.6	9.0	0.2	8.0	88.9	8	0.2	7	87.5

1 Average, rounded to one decimal place.

2 Rate per 100 000 population based on 2006 census data.

3 Proportion (%) of total cases who were confirmed.

Table 2. Age distribution of total and confirmed cases of meningococcal disease and proportion of confirmed to total cases, January 2005/2006 (average), 2007/2008 (average) and 2009

Age Group	Jan 2005/2006 average				Jan 2007/2008 average				Jan 2009			
	No. Total ¹	Rate ²	No. Conf ¹	% ³	No. Total ¹	Rate ²	No. Conf ¹	% ³	No. Total	Rate ²	No. Conf	% ³
<1	2.0	3.5	1.0	50.0	0.0	0.0	0.0	-	2	3.5	2	100
1-4	3.5	1.6	3.0	85.7	3.5	1.6	3.0	85.7	4	1.8	3	75.0
5-9	1.5	0.5	1.0	66.7	1.5	0.5	1.0	66.7	0	0.0	0	-
10-14	2.0	0.7	1.5	75.0	0.0	0.0	0.0	-	1	0.3	1	100
15-19	2.0	0.7	2.0	100	1.0	0.3	1.0	100	0	0.0	0	-
20-29	2.0	0.4	2.0	100	1.0	0.2	1.0	100	0	0.0	0	-
30-39	1.0	0.2	0.5	50.0	0.5	0.1	0.5	100	0	0.0	0	-
40+	1.5	0.1	1.5	100	1.5	0.1	1.5	100	1	0.1	1	100
Total	15.5	0.4	12.5	80.6	9.0	0.2	8.0	88.9	8	0.2	7	87.5

1 Average, rounded to one decimal place.

2 Rate per 100 000 population based on 2006 census data.

3 Proportion (%) of total cases who were confirmed.

Table 3. Meningococcal disease cases less than two years of age, January 2005/2006 (average), 2007/2008 (average) and 2009

Age (months)	Jan 2005/2006 average				Jan 2007/2008 average				Jan 2009			
	Total cases		Confirmed cases		Total cases		Confirmed cases		Total cases		Confirmed cases	
	No. ¹	Cum % ² (n=15.5)	No. ¹	Cum % ² (n=12.5)	No. ¹	Cum % ² (n=9.0)	No. ¹	Cum % ² (n=8.0)	No.	Cum % ² (n=8)	No.	Cum % ² (n=7)
<1	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0	0.0	0	0.0
1	0.5	3.2	0.5	4.0	0.0	0.0	0.0	0.0	1	12.5	1	14.3
2	1.0	9.7	0.0	4.0	0.0	0.0	0.0	0.0	0	12.5	0	14.3
3	0.5	12.9	0.5	8.0	0.0	0.0	0.0	0.0	0	12.5	0	14.3
4	0.0	12.9	0.0	8.0	0.0	0.0	0.0	0.0	0	12.5	0	14.3
5	0.0	12.9	0.0	8.0	0.0	0.0	0.0	0.0	0	12.5	0	14.3
6	0.0	12.9	0.0	8.0	0.0	0.0	0.0	0.0	0	12.5	0	14.3
7	0.0	12.9	0.0	8.0	0.0	0.0	0.0	0.0	0	12.5	0	14.3
8	0.0	12.9	0.0	8.0	0.0	0.0	0.0	0.0	0	12.5	0	14.3
9	0.0	12.9	0.0	8.0	0.0	0.0	0.0	0.0	0	12.5	0	14.3
10	0.0	12.9	0.0	8.0	0.0	0.0	0.0	0.0	0	12.5	0	14.3
11	0.0	12.9	0.0	8.0	0.0	0.0	0.0	0.0	1	25.0	1	28.6
12	0.5	16.1	0.0	8.0	0.0	0.0	0.0	0.0	0	25.0	0	28.6
13	0.0	16.1	0.0	8.0	1.0	11.1	0.5	6.3	0	25.0	0	28.6
14	0.5	19.4	0.5	12.0	0.0	11.1	0.0	6.3	0	25.0	0	28.6
15	0.0	19.4	0.0	12.0	0.5	16.7	0.5	12.5	0	25.0	0	28.6
16	0.0	19.4	0.0	12.0	0.5	22.2	0.5	18.8	0	25.0	0	28.6
17	0.0	19.4	0.0	12.0	0.0	22.2	0.0	18.8	0	25.0	0	28.6
18	0.0	19.4	0.0	12.0	0.0	22.2	0.0	18.8	0	25.0	0	28.6
19	0.0	19.4	0.0	12.0	0.0	22.2	0.0	18.8	0	25.0	0	28.6
20	0.0	19.4	0.0	12.0	0.0	22.2	0.0	18.8	0	25.0	0	28.6
21	0.0	19.4	0.0	12.0	0.5	27.8	0.5	25.0	0	25.0	0	28.6
22	0.0	19.4	0.0	12.0	0.0	27.8	0.0	25.0	0	25.0	0	28.6
23	0.0	19.4	0.0	12.0	0.0	27.8	0.0	25.0	0	25.0	0	28.6
Total	3.0		1.5		2.5		2.0		2		2	

1 Average, rounded to one decimal place.

2 Cumulative percentage of cases in all age groups.

Table 4. Meningococcal disease cases by District Health Board and age group, January 2005/2006 (average), 2007/2008 (average), 2007/2008 (average) and 2009

District Health Board	Jan 2005/2006 average										Jan 2007/2008 average										Jan 2009					
	<1	1-4	5-9	10-14	15-19	20-29	30+	Total	<1	1-4	5-9	10-14	15-19	20-29	30+	Total	<1	1-4	5-9	10-14	15-19	20-29	30+	Total		
Northland	0.0	0.5	0.0	0.0	0.0	0.0	0.0	0.5	0.0	0.5	0.0	0.0	0.0	0.0	0.0	0.5	0	0	0	0	0	0	0	0		
Waitemata	1.0	0.5	0.0	0.0	0.5	1.0	0.5	3.5	0.0	0.5	0.0	0.0	0.5	0.0	0.0	1.0	0	0	0	0	0	0	0	0		
Auckland	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.5	0.0	0.5	0	0	0	0	0	0	0	0		
Counties Manukau	0.0	0.0	0.5	0.0	0.5	0.0	0.0	1.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	1	0	0	1	0	0	0	2		
Waikato	0.0	0.0	0.0	0.0	0.0	0.0	0.5	0.5	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0	0	0	0	0	0	0	0		
Lakes	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0	0	0	0	0	0	0	0		
Bay of Plenty	0.0	0.0	0.5	0.0	0.0	0.0	0.0	0.5	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0	1	0	0	0	0	0	1		
Tarawhiti	0.0	0.5	0.0	0.0	0.0	0.0	0.0	0.5	0.0	0.5	0.0	0.0	0.0	0.0	0.0	0.5	0	0	0	0	0	0	0	0		
Taranaki	0.5	0.0	0.0	0.0	0.5	0.0	0.0	1.0	0.0	0.5	0.0	0.0	0.0	0.0	0.0	0.5	0	0	0	0	0	0	0	0		
Hawke's Bay	0.0	1.0	0.0	0.0	0.0	0.0	0.0	1.0	0.0	1.0	0.5	0.0	0.0	0.0	0.5	2.0	0	1	0	0	0	0	0	1		
Whanganui	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0	0	0	0	0	0	0	0		
MidCentral	0.0	0.0	0.0	1.0	0.0	0.0	0.0	1.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0	0	0	0	0	0	0	0		
Hutt	0.0	0.0	0.0	0.0	0.0	0.0	0.5	0.5	0.0	0.0	0.0	0.0	0.0	0.0	0.5	0.5	0	0	0	0	0	0	0	0		
Capital and Coast	0.0	0.5	0.5	0.0	0.0	0.0	0.0	1.0	0.0	0.0	0.0	0.0	0.5	0.0	0.0	0.5	1	0	0	0	0	0	1	2		
Wairarapa	0.0	0.0	0.0	0.5	0.0	0.0	0.0	0.5	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0	0	0	0	0	0	0	0		
Nelson																										
Marlborough	0.0	0.0	0.0	0.0	0.0	1.0	0.0	1.0	0.0	0.0	0.5	0.0	0.0	0.0	0.0	0.5	0	0	0	0	0	0	0	0		
West Coast	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0	0	0	0	0	0	0	0		
Canterbury	0.0	0.5	0.0	0.0	0.5	0.0	1.0	2.0	0.0	0.5	0.5	0.0	0.0	0.5	0.0	1.5	0	1	0	0	0	0	0	1		
South Canterbury	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0	0	0	0	0	0	0	0		
Otago	0.5	0.0	0.0	0.0	0.0	0.0	0.0	0.5	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0	1	0	0	0	0	0	1		
Southland	0.0	0.0	0.0	0.5	0.0	0.0	0.0	0.5	0.0	0.0	0.0	0.0	0.0	0.0	1.0	1.0	0	0	0	0	0	0	0	0		
NZ Total	2.0	3.5	1.5	2.0	2.0	2.0	2.5	15.5	0.0	3.5	1.5	0.0	1.0	1.0	2.0	9.0	2	4	0	1	0	0	1	8		

1 Average, rounded to one decimal place.

Table 5. Numbers and age-standardised incidence rates by ethnicity for cases of meningococcal disease and proportion of confirmed to total cases, January 2005/2006 (average), 2007/2008 (average) and 2009

Ethnicity	Jan 2005/2006 average				Jan 2007/2008 average				Jan 2009			
	No. Total ¹	Cum Rate ²	No. Conf ¹	% ³	No. Total ¹	Cum Rate ²	No. Conf ¹	% ³	No. Total	Cum Rate ²	No. Conf	% ³
European	9.0	0.4	7.0	77.8	4.5	0.2	3.5	77.8	2	0.1	1	50.0
Maori ⁴	3.5	0.4	3.5	100	2.0	0.2	2.0	100	3	0.3	3	100
Pacific ⁵	1.5	0.4	0.5	33.3	1.0	0.3	1.0	100	3	1.3	3	100
Other	0.5	0.1	0.5	100	1.0	0.3	1.0	100	0	0.0	0	-
Unknown	1.0		1.0	100	0.5		0.5	100	0		0	-
Total	15.5	0.4	12.5	80.6	9.0	0.2	8.0	88.9	8	0.2	7	87.5

1 Average, rounded to one decimal place.

2 Cumulative rate per 100 000 direct standardised to age distribution of the total NZ population (based on 2006 census data).

3 Proportion (%) of total cases who were confirmed.

4 Rate is calculated using mixed Maori ethnicity.

5 Rate is calculated using mixed Pacific people ethnicity (excluding Maori).

Table 6. Meningococcal disease cases by strain type, January 2005/2006 (average), 2007/2008 (average) and 2009

Strain Type	Jan 2005/2006 average		Jan 2007/2008 average		Jan 2009	
	No. ¹	% ²	No. ¹	% ²	No.	% ²
Epidemic Strain	8.5	70.8	4.0	53.3	4	66.7
B Other	1.0	8.3	3.5	46.7	1	16.7
C	0.5	4.2	0.0	0.0	1	16.7
W, Y, Z	2.0	16.7	0.0	0.0	0	0.0
Total with determined strain	12.0	100	7.5	100	6	100
Confirmed but awaiting strain result	0.5		0.0		1	
DNA positive, type not determined	0.0		0.5		0	
Probable	3.0		1.0		1	
Total with undetermined strain	3.5		1.5		2	
Grand total	15.5		9.0		8	

1 Average, rounded to one decimal place.

2 Proportion (%) of cases where a strain has been determined.

Table 7. Meningococcal disease cases by strain type, February-January: 2007, 2008 and 2009

Strain Type	Feb 06-Jan 07		Feb 07-Jan 08		Feb 08-Jan 09	
	No.	Rate ¹	No.	Rate ¹	No.	Rate ¹
Epidemic Strain	77	1.9	45	1.1	45	1.1
B Other	26	0.6	30	0.7	33	0.8
C	24	0.6	9	0.2	10	0.2
W, Y, Z	8	0.2	6	0.1	11	0.3
Total with defined strain	135	3.4	90	2.2	99	2.5
Total confirmed with strain undetermined	10	0.2	5	0.1	9	0.2
Grand total	145	3.6	95	2.4	108	2.7

¹ Cumulative rate per 100 000 based on 2006 census data.

Table 8. Epidemic strain meningococcal disease by region, February-January: 2007, 2008 and 2009

Region	Feb 06-Jan 07		Feb 07-Jan 08		Feb 08-Jan 09	
	No.	Rate ¹	No.	Rate ¹	No.	Rate ¹
Northern	34	2.3	17	1.2	12	0.8
Midland	18	2.3	9	1.2	9	1.2
Central	12	1.5	12	1.5	16	2.0
Southern	13	1.3	7	0.7	8	0.8
Grand total	77	1.9	45	1.1	45	1.1

¹ Cumulative rate per 100 000 based on 2006 census data.

Table 9. Epidemic strain meningococcal disease by age group, February-January: 2007, 2008 and 2009

Age group (years)	Feb 06-Jan 07		Feb 07-Jan 08		Feb 08-Jan 09	
	No.	Rate ¹	No.	Rate ¹	No.	Rate ¹
0-4	31	11.3	25	9.1	25	9.1
5-19	17	1.9	13	1.5	7	0.8
20+	29	1.0	7	0.2	13	0.5
Grand total	77	1.9	45	1.1	45	1.1

¹ Cumulative rate per 100 000 based on 2006 census data.

Table 10. Epidemic strain meningococcal disease by ethnicity, February-January: 2007, 2008 and 2009

Ethnicity	Feb 06-Jan 07		Feb 07-Jan 08		Feb 08-Jan 09	
	No.	Rate ¹	No.	Rate ¹	No.	Rate ¹
European	37	1.5	15	0.6	13	0.5
Maori ²	25	2.9	19	2.3	17	1.9
Pacific Peoples ³	13	5.2	9	2.6	10	3.8
Other	2	0.4	1	0.3	3	0.7
Unknown	0		1		2	
Grand total	77	1.9	45	1.1	45	1.1

¹ Cumulative rate per 100 000 direct standardised to age distribution of the total NZ population (based on 2006 census data).

² Rate is calculated using mixed Maori ethnicity.

³ Rate is calculated using mixed Pacific people ethnicity (excluding Maori).

Table 11. Meningococcal disease cases by District Health Board and strain type, January 2005/2006 (average), 2007/2008 (average) and 2009

District Health Board	Jan 2005/2006 average			Jan 2007/2008 average			Jan 2009							
	Epidemic Strain		Other ¹	Epidemic Strain		Other ¹	Epidemic Strain		Other ¹					
	No. ²	% ³		No. ²	% ³		No. ²	% ³		No. ²	% ³			
Northland	0.5	100	0.0	0.0	0.5	100	0.0	0.0	0	-	0	-		
Waitemata	1.5	100	0.0	0.0	0.5	50.0	0.0	50.0	0	-	0	-		
Auckland	0.0	-	0.0	-	0.5	100	0.0	0.0	0	-	0	-		
Counties Manukau	0.5	50.0	0.5	50.0	0.0	0.0	0.0	0.0	0	0.0	1	50.0		
Waikato	0.5	100	0.0	0.0	0.0	0.0	0.0	0.0	0	-	0	-		
Lakes	0.0	-	0.0	-	0.0	0.0	0.0	0.0	0	-	0	-		
Bay of Plenty	0.5	100	0.0	0.0	0.0	0.0	0.0	0.0	1	100	0	0.0		
Tarawhiti	0.5	100	0.0	0.0	0.0	0.0	0.0	0.0	0	-	0	-		
Taranaki	0.5	50.0	0.0	0.0	0.5	50.0	0.0	0.0	0	-	0	-		
Hawke's Bay	0.5	50.0	0.0	0.0	0.5	50.0	1.5	100	0.0	0.0	0	-		
Whanganui	0.0	-	0.0	-	0.0	0.0	0.0	0.0	0	-	0	-		
MidCentral	1.0	100	0.0	0.0	0.0	0.0	0.0	0.0	0	-	0	-		
Hutt	0.0	0.0	0.0	0.0	0.5	100	0.0	0.0	0.5	100	0	-		
Capital and Coast	0.5	100	0.0	0.0	0.0	0.0	0.0	0.0	0	-	2	100		
Wairarapa	0.0	-	0.0	-	0.0	0.0	0.0	0.0	0	-	0	-		
Nelson	0.5	50.0	0.0	0.0	0.5	50.0	0.0	0.0	0.5	100	0	-		
Marlborough	0.0	-	0.0	-	0.0	0.0	0.0	0.0	0	-	0	-		
West Coast	1.0	66.7	0.0	0.0	0.5	33.3	0.5	50.0	0.0	0.0	0	-		
Canterbury	0.0	-	0.0	-	0.0	0.0	0.0	0.0	0	-	0	-		
South Canterbury	0.0	0.0	0.0	0.0	0.5	100	0.0	0.0	1	100	0	0.0		
Otago	0.5	100	0.0	0.0	0.0	0.0	0.5	50.0	0.0	0.0	0	-		
Southland	0.5	100	0.0	0.0	0.0	0.0	0.0	0.0	0.5	50.0	0	-		
New Zealand total	8.5	70.8	0.5	4.2	3.0	25.0	4.0	53.3	0.0	0.0	4	66.7	1	16.7

1 Other includes: non epidemic group B, W, Y, Z.

2 Average, rounded to one decimal place.

3 Proportion (%) of cases where a strain type has been determined for each DHB.

Table 12. Meningococcal disease cases by age group and strain type, January 2005/2006 (average), 2007/2008 (average) and 2009

Age Group	Jan 2005/2006 average						Jan 2007/2008 average						Jan 2009					
	Epidemic Strain		Group C		Other ¹		Epidemic Strain		Group C		Other ¹		Epidemic Strain		Group C		Other ¹	
	No. ²	% ³	No. ²	% ³	No. ²	% ³	No. ²	% ³	No. ²	% ³	No. ²	% ³	No. ²	% ³	No. ²	% ³	No. ²	% ³
<1	0.0	0.0	0.0	0.0	1.0	100	0.0	-	0.0	-	0.0	-	1	50.0	0	0.0	1	50.0
1-4	2.5	83.3	0.0	0.0	0.5	16.7	1.0	33.3	0.0	0.0	2.0	66.7	2	100	0	0.0	0	0.0
5-9	0.5	50.0	0.5	50.0	0.0	0.0	0.5	50.0	0.0	0.0	0.5	50.0	0	-	0	-	0	-
10-14	1.5	100	0.0	0.0	0.0	0.0	0.0	-	0.0	-	0.0	-	0	0.0	1	100	0	0.0
15-19	1.5	100	0.0	0.0	0.0	0.0	0.5	100	0.0	0.0	0.0	0.0	0	-	0	-	0	-
20-29	1.5	75.0	0.0	0.0	0.5	25.0	1.0	100	0.0	0.0	0.0	0.0	0	-	0	-	0	-
30-39	0.5	100	0.0	0.0	0.0	0.0	0.5	100	0.0	0.0	0.0	0.0	0	-	0	-	0	-
40+	0.5	33.3	0.0	0.0	1.0	66.7	0.5	33.3	0.0	0.0	1.0	66.7	1	100	0	0.0	0	0.0
Total	8.5	70.8	0.5	4.2	3.0	25.0	4.0	53.3	0.0	0.0	3.5	46.7	4	66.7	1	16.7	1	16.7

1 Other includes: non epidemic group B, W, Y, Z.

2 Average, rounded to one decimal place.

3 Proportion (%) of cases where a strain type has been determined for each DHB.

Table 13. Meningococcal disease cases by basis for diagnosis, January 2005/2006 (average), 2007/2008 (average) and 2009

Basis for diagnosis ¹	Jan 2005/2006 average		Jan 2007/2008 average		Jan 2009	
	No. ²	% ³	No. ²	% ³	No.	% ³
Isolation of <i>N. meningitidis</i> from blood and/or CSF or any other sterile site	10.5	67.7	6.5	72.2	5	62.5
PCR test	2.0	12.9	1.5	16.7	2	25.0
Gram-negative diplococci in CSF	0.0	0.0	0.0	0.0	0	0.0
Confirmed –subtotal	12.5	80.6	8.0	88.9	7	87.5
Clinical criteria and a positive throat swab	0.0	0.0	0.0	0.0	0	0.0
Clinical criteria	3.0	19.4	1.0	11.1	1	12.5
Probable –subtotal	3.0	19.4	1.0	11.1	1	12.5
Total	15.5	100	9.0	100	8	100

1 Each case is represented only once in the table.

2 Average, rounded to one decimal place.

3 Proportion (%) of total cases.

Table 14. Case-fatality rates for meningococcal disease cases by District Health Board, age, ethnicity, and group, January 2005/2006 (average), 2007/2008 (average) and 2009

Features of case and infecting organism	Jan 2005/2006 average		Jan 2007/2008 average		Jan 2009	
	No. fatalities ¹	Case fatality rate (%)	No. Fatalities ¹	Case fatality rate (%)	No. fatalities	Case fatality rate (%)
Northland	0.0	0.0	0.0	0.0	0	-
Waitemata	0.0	0.0	0.0	0.0	0	-
Auckland	0.0	-	0.0	0.0	0	-
Counties Manukau	0.5	50.0	0.0	-	0	0.0
Waikato	0.0	0.0	0.0	-	0	-
Lakes	0.0	-	0.0	-	0	-
Bay of Plenty	0.0	0.0	0.0	-	0	0.0
Tairāwhiti	0.0	0.0	0.0	0.0	0	-
Taranaki	0.0	0.0	0.0	0.0	0	-
Hawke's Bay	0.0	0.0	0.5	25.0	0	0.0
Whanganui	0.0	-	0.0	-	0	-
MidCentral	0.0	0.0	0.0	-	0	-
Hutt	0.0	0.0	0.0	0.0	0	-
Capital and Coast	0.0	0.0	0.0	0.0	0	0.0
Wairarapa	0.0	0.0	0.0	-	0	-
Nelson Marlborough	0.0	0.0	0.0	0.0	0	-
West Coast	0.0	-	0.0	-	0	-
Canterbury	0.5	25.0	0.5	33.3	0	0.0
South Canterbury	0.0	-	0.0	-	0	-
Otago	0.0	0.0	0.0	-	0	0.0
Southland	0.0	0.0	0.0	0.0	0	-
<1 year	0.0	0.0	0.0	-	0	0.0
1-4 years	0.0	0.0	0.5	14.3	0	0.0
5-9 years	0.5	33.3	0.5	33.3	0	-
10-14 years	0.0	0.0	0.0	-	0	0.0
15-19 years	0.0	0.0	0.0	0.0	0	-
20-29 years	0.0	0.0	0.0	0.0	0	-
30-39 years	0.5	50.0	0.0	0.0	0	-
40+ years	0.0	0.0	0.0	0.0	0	0.0
European	0.0	0.0	1.0	22.2	0	0.0
Māori	0.5	14.3	0.0	0.0	0	0.0
Pacific	0.0	0.0	0.0	0.0	0	0.0
Other	0.0	0.0	0.0	0.0	0	-
Unknown	0.5	50.0	0.0	0.0	0	-
Epidemic Strain	0.5	5.9	0.5	12.5	0	0.0
B Other	0.0	0.0	0.0	0.0	0	0.0
C	0.5	100	0.0	-	0	0.0
Other	0.0	0.0	0.0	-	0	-
Confirmed but awaiting strain result/not done	0.0	0.0	0.0	-	0	0.0
Confirmed by gramneg - no organism	0.0	-	0.0	-	0	-
DNA positive, type not determined	0.0	-	0.0	0.0	0	-
Probable	0.0	0.0	0.5	50.0	0	0.0
Total	1.0	6.5	1.0	11.1	0	0.0

¹ Average, rounded to one decimal place.

REFERENCES

1. Ministry of Health, *Communicable Disease Control Manual*. 1998.
2. Bennett D. E. and Cafferkey M. T, *Consecutive use of two multiplex PCR-based assays for simultaneous identification and determination of capsular status of nine common Neisseria meningitidis serogroups associated with invasive disease*. J Clin Microbiol, 2006. 44: p. 1127-31.
3. Abdillahi H and Poolman JT, *Whole-cell ELISA for typing Neisseria meningitidis with monoclonal antibodies*. FEMS Microbiol Lett, 1987. 48: p. 367 - 371.
4. Saunders NB, Zollinger WD, and Rao VB, *A rapid and sensitive PCR strategy employed for amplification and sequencing of the porA gene of Neisseria meningitidis*. Gene, 1993. 137: p. 153-162.
5. Maiden M. C., et al., *Multilocus sequence typing: a portable approach to the identification of clones within populations of pathogenic microorganisms*. Proc Natl Acad Sci USA, 1998. 95: p. 3140-5.
6. Bygraves J. A. and Maiden M. C. J., *Analysis of the clonal relationships between strains of Neisseria meningitidis by pulsed field gel electrophoresis*. J Gen Microbiol, 1992. 138: p. 523-531.