

# PERTUSSIS REPORT

---

This report includes cases of pertussis reported in EpiSurv up to midnight 23 November 2012. Data were extracted from EpiSurv at 10.00 am 27 November 2012.

## Summary

In the past two surveillance weeks (10 November – 23 November 2012), 292 new cases of pertussis (133 and 159 cases, respectively) were notified, including 145 confirmed cases, 81 probable cases, 8 suspect cases, and 58 cases still under investigation. Fewer cases were reported in the past two surveillance weeks compared to the numbers reported over the previous two weeks (330 cases). Twenty-six (8.9%) of the notified cases were aged less than 1 year. Nineteen cases were hospitalised.

There has been a total of 5267 pertussis notifications reported in EpiSurv since the first surveillance week of 2012 (compared to 1392 over the same period in 2011), including 2177 confirmed cases, 2657 probable cases, 257 suspect cases, and 176 cases still under investigation. 362 (6.9%) of the notified cases were in the less than 1 year age group. During this period, 269 hospitalisations and two deaths have been reported.

In the last two weeks, the highest number of cases (excluding cases under investigation) was reported in Canterbury (55 cases), Waikato and Capital and Coast (22 cases each), and MidCentral (21 cases) DHBs. The highest cumulative rate to date in 2012 was recorded in Nelson Marlborough (425.3 per 100 000, 595 cases), followed by West Coast (400.5 per 100 000, 132 cases) and Tairāwhiti (255.4 per 100 000, 119 cases) DHBs. During this same period the highest number of notifications was reported from Canterbury DHB (1111 cases), followed by Capital and Coast (623) and Nelson Marlborough (595), Hutt Valley (302) and Counties Manukau (292) DHBs. Monthly pertussis rates and cases (excluding cases under investigation) by DHB can be seen in Figures 8 and 9 (appendix).

This report summarises pertussis notifications for 2012 (first surveillance week starts on 31 December 2011) and new cases in the last two weeks, and incorporates the temporal distribution of cases, the distribution of cases by age, ethnicity (prioritised), and DHB, as well as hospitalisations and immunisation status. The case classification used in this report is specified in the appendix. Case definitions have changed following the release of the Ministry of Health's *Communicable Disease Control Manual 2012* on 31 May 2012.

## Temporal distribution of pertussis cases

Figure 1 shows weekly total pertussis notifications for 2010, 2011 and 2012 (to week ending 23 November). Notifications for the past two weeks of 2012 remain well above 2011 and 2010 levels. Since week 34 in 2011 (ending 26 August) notifications have been increasing more or less consistently. The highest weekly notification count occurred during week 44 of this year. Note the total number of notifications may change as cases are investigated further and some are found not to meet the case definition. Two deaths have been reported since the beginning of this year. Figure 5 (appendix) shows weekly pertussis notifications for confirmed, suspect and probable cases only for 2010, 2011 and 2012.

**Figure 1: Number of pertussis notifications by week reported 2010 - 2012**





Figure 2 shows pertussis notifications and hospitalisations by calendar month, and notifications in those aged less than 1 year between 1 January 1997 and 31 October 2012. A four to five-year cycle can be seen with large peaks in notifications in years 2000 and 2004 and a much smaller peak in 2009. However, notifications have been rising again since August 2011. Increases in hospitalisations show a similar cycle, although peaks in hospitalisations do not always coincide with peaks in notifications. Figure 6 (appendix) shows annual rates in the less than 1 year age group over 1997-2011.

**Figure 2: Pertussis notifications and hospitalisations by calendar month-year since 1997 up to 31 October 2012**

In the following sections all analyses exclude cases still under investigation. Therefore, “cases” refer to those classified as confirmed, probable, or suspect.

## Age distribution of cases

Table 1 shows notifications and associated rates by age, including new cases for the last two weeks. Pertussis rates varied across age groups. Of the cases reported in 2012, infants aged less than one year had the highest cumulative notification rate (558.0 per 100 000 population, 348 cases), followed by the 1 to 4 years (320.8 per 100 000, 808 cases), and 5 to 9 years (229.8 per 100 000, 660 cases) age groups.

Of the 5090 cumulative cases with known age, 40 (0.8%) were infants under 6 weeks of age. Figure 3 shows the cumulative notification rate of pertussis cases by age group and ethnicity in 2012.

**Table 1: Pertussis cases and rates by age group in 2012, and new cases in the last two weeks**

Age group (Years)	Cumulative <sup>2</sup> notifications			Last two weeks <sup>3</sup>	
	All cases <sup>1</sup>	Rates <sup>1</sup>	Hosp	New Cases	Hosp
<1	348	558.0	149	21	8
1 to 4	808	320.8	26	26	3
5 to 9	660	229.8	9	33	0
10 to 14	474	161.8	5	21	0
15 to 19	242	76.3	6	7	1
20 to 29	447	72.2	4	14	0
30 to 39	590	104.8	10	34	1
40 to 49	647	102.4	13	28	1
50 to 59	406	73.0	14	30	1
60 to 69	289	69.3	12	9	1
70+	179	44.0	13	10	1
Unknown	1		0	1	0
<b>Overall</b>	<b>5091</b>	<b>115.6</b>	<b>261</b>	<b>234</b>	<b>17</b>

<sup>1</sup>Rate of pertussis cases per 100 000 population calculated using 2011 mid-year population estimates.

<sup>2</sup>Cumulative notifications (excluding cases under investigation) since 31 December 2011

<sup>3</sup>Notifications between 10 November and 23 November 2012 inclusive

**Hosp:** hospitalisation counts

## Ethnicity

Pertussis cases and rates by ethnicity are shown in Table 2. Of the pertussis cases with known ethnicity, the European ethnic group had the highest numbers reported in the last two weeks (177 cases). Of the cases in 2012, the ethnic-specific cumulative rates were highest for the European ethnic group (141.5 per 100 000, 3813 cases), followed by Māori (120.1 per 100 000, 679 cases) and Pacific Peoples (77.3 per 100 000, 175 cases). Figure 3 shows the European ethnic group having the highest notification rates across all age groups except the under 1 year age group. The ethnic distribution of cases in the under 1 year age group is also shown below. Māori had the highest notification rates in this age group, followed by Pacific Peoples.

**Table 2: Pertussis cases and rates by ethnicity (prioritised) in 2012, and new cases in the last two weeks**

Ethnicity	Cumulative <sup>2</sup> notifications					Last two weeks <sup>3</sup>		
	All cases (Rate <sup>1</sup> )		<1 year (Rate <sup>1</sup> )		Hosp (% <sup>4</sup> )	New Cases	<1 year	Hosp
Māori	679	(120.1)	111	(791.1)	86 (12.7)	27	7	5
Pacific Peoples	175	(77.3)	35	(683.5)	37 (21.1)	6	1	0
Other	175	(46.7)	13	(247.5)	12 (6.9)	4	0	1
European	3813	(141.5)	179	(602.6)	121 (3.2)	177	11	9
Unknown	249		10		5	20	2	2
<b>Overall</b>	<b>5091</b>	<b>(126.4)</b>	<b>348</b>	<b>(614.5)</b>	<b>261 (5.1)</b>	<b>234</b>	<b>21</b>	<b>17</b>


<sup>1</sup>Value in brackets denotes rate of pertussis cases per 100 000 population calculated using usually resident populations (Census 2006).

<sup>2</sup>Cumulative notifications (excluding cases under investigation) since 31 December 2011

<sup>3</sup>Notifications between 10 November and 23 November 2012 inclusive

<sup>4</sup>Percentage of hospitalised notifications by ethnic group

**Figure 3: Pertussis rates per 100 000 populations by age group and ethnicity, in 2012**



**Note:** Rate of pertussis cases per 100 000 population calculated using Census 2006 usually resident populations.

Figure 7 (appendix) shows the trend of cumulative pertussis notification rates (per 100 000 population) by age group and ethnicity for years 2003 to 2011. Over this time period rates have been generally highest among Pacific Peoples in the less than 1 year age group, while

in other age groups rates have been consistently highest in the European ethnic group. Note that these rates are for all notifications.


## Hospitalisations and Deaths

The distribution of hospitalisations by age group, ethnicity, and DHB is described in Table 1, Table 2 and Table 5, respectively. In the last two weeks, 17 hospitalisations were recorded. There have been 261 hospitalisations reported in EpiSurv in 2012. One hundred and forty-nine (57.1%) of these were infants aged less than one year including 36 cases aged less than six weeks. Of the 4457 cases with known ethnicity and hospitalisation status, the ethnic-specific proportions of hospitalisations were as follows: Pacific Peoples (23.0%, 37/161), Māori (13.8%, 86/621), Other (7.6%, 12/158), and European (3.4%, 121/3517). There have been two pertussis deaths reported in 2012. One in a 3 year old unimmunised child with underlying health conditions, and one in an infant aged less than 6 weeks with a history of premature birth.

## Geographic distribution

The rates of pertussis notifications by DHB can be seen in Figure 4 and Table 5 (appendix). In the last two weeks, the highest number of cases was reported in Canterbury (55 cases), Waikato and Capital and Coast (22 cases each), and MidCentral (21 cases) DHBs. The highest cumulative rate in 2012 was recorded in Nelson Marlborough (425.3 per 100 000, 595 cases), followed by West Coast (400.5 per 100 000, 132 cases) and Tairāwhiti (255.4 per 100 000, 119 cases) DHBs. The highest number of notifications was reported from Canterbury DHB (1111 cases), followed by Capital and Coast (623), Nelson Marlborough (595), Hutt Valley (302) and Counties Manukau (292) DHBs. Cases in the under 1 year age group by DHB are shown in Table 5 (appendix). Also, monthly pertussis rates and cases (excluding cases under investigation) by DHB can be seen in Figures 8 and 9 (appendix).

**Figure 4: Pertussis notifications rates (cases per 100 000 populations) in 2012**



**Note:** Rates were calculated using 2011 mid-year population estimates.

## Immunisation status

The immunisation status for confirmed pertussis cases with known age is shown in Table 3 and Table 4 for the last two weeks and for 2012, respectively. Of the 145 confirmed cases reported in the last two weeks, 85 (58.6%) had a known vaccination status. Of these 85 cases, 29 were not vaccinated. Five cases had received one dose of vaccine, five cases had received two doses, 12 cases had received three doses, 14 cases had received four doses, and eight reported having completed pertussis vaccination. A further 12 cases reported being vaccinated but no dose information was available.

**Table 3: Immunisation status of pertussis cases (confirmed) notified in the last two weeks (ending 23 November)**

Age Group	Total cases	One dose	Two doses	Three doses	Four doses	Five doses	Vaccinated		
							(no dose info)	Not vaccinated	Unknown
<6wks	2	0	0	0	0	0	0	2	0
6wks - 2mths	7	3	0	0	0	0	0	3	1
3-4 mths	4	1	3	0	0	0	0	0	0
5mths - 3yrs	17	0	0	10	1	0	0	2	4
4 - 10yrs	33	0	0	1	13	6	2	5	6
11+ yrs	81	1	2	1	0	2	9	17	49
Unknown	1	0	0	0	0	0	1	0	0
<b>Total</b>	<b>145</b>	<b>5</b>	<b>5</b>	<b>12</b>	<b>14</b>	<b>8</b>	<b>12</b>	<b>29</b>	<b>60</b>

**Note:** Immunisation status has been extracted from Episurv notifications. Health professionals use a range of sources to update immunisation status including the NIR, parental recall or Well Child book records.

Of the 2176 confirmed cases with known age reported during 2012, 1433 (65.9%) had a known vaccination status (Table 4). Of these 1433 cases, 517 were not vaccinated, including 22 cases aged less than 6 weeks and thus not eligible for vaccination. One hundred and thirteen cases had received one dose of vaccine, 38 cases had received two doses, 198 cases had received three doses, 188 cases had received four doses, and 112 cases reported having completed pertussis vaccination. A further 268 cases reported being vaccinated but no dose information was available.

**Table 4: Immunisation status of pertussis cases (confirmed) notified in 2012 (since 31 December 2011)**

Age Group	Total cases	One dose	Two doses	Three doses	Four doses	Five doses	Vaccinated		
							(no dose info)	Not vaccinated	Unknown
<6wks	25	0	0	0	0	0	0	22	3
6wks - 2mths	76	41	1	0	0	0	3	26	5
3-4 mths	39	7	17	0	0	0	0	14	1
5mths - 3yrs	355	5	10	151	31	1	19	105	33
4 - 10yrs	542	9	4	33	136	73	74	156	57
11+ yrs	1139	51	6	14	21	38	171	194	644
Unknown	1	0	0	0	0	0	1	0	0
<b>Total</b>	<b>2177</b>	<b>113</b>	<b>38</b>	<b>198</b>	<b>188</b>	<b>112</b>	<b>268</b>	<b>517</b>	<b>743</b>

**Note:** Immunisation status has been extracted from Episurv notifications. Health professionals use a range of sources to update immunisation status including the NIR, parental recall or Well Child book records.

## Appendix

**Table 5: Pertussis cases and rates by DHB in 2012, and new cases in the last two weeks**

DHB	Cumulative <sup>2</sup> notifications				Last two weeks <sup>3</sup>		
	All cases	Rates <sup>1</sup>	<1 year*	Hosp	New Cases	<1 year*	Hosp
Northland	72	45.5	7	4	2	1	0
Waitemata	222	40.7	19	21	7	1	3
Auckland	234	51.2	20	24	5	1	0
Counties Manukau	292	58.4	40	51	12	1	5
Waikato	282	76.7	23	17	22	5	3
Lakes	89	86.4	13	11	1	0	0
Bay of Plenty	148	69.8	7	3	16	0	0
Tairāwhiti	119	255.4	14	4	1	0	0
Taranaki	116	105.6	5	7	10	0	0
Hawke's Bay	122	78.3	12	6	4	1	0
Whanganui	139	220.4	17	12	4	1	0
MidCentral	181	107.5	16	9	21	4	1
Hutt Valley	302	209.0	12	5	18	0	0
Capital and Coast	623	211.4	29	9	22	0	0
Wairarapa	84	207.0	6	11	10	1	1
Nelson Marlborough	595	425.3	33	8	12	1	0
West Coast	132	400.5	3	2	2	0	0
Canterbury	1111	221.0	49	35	55	4	3
South Canterbury	26	46.1	0	2	0	0	0
Southern	202	65.9	23	20	10	0	1
<b>Total</b>	<b>5091</b>	<b>115.6</b>	<b>348</b>	<b>261</b>	<b>234</b>	<b>21</b>	<b>17</b>


<sup>1</sup>Rate of pertussis cases per 100 000 population calculated using 2011 mid-year population estimates.

<sup>2</sup>Cumulative notifications (excluding cases under investigation) since 31 December 2011

<sup>3</sup>Notifications between 10 November and 23 November 2012 inclusive


\*Cases in the less than 1 year age group

**Figure 5: Comparative trend of pertussis cases by week reported during years 2010, 2011 and 2012**




**Note:** Includes confirmed, probable and suspect cases only.

Figure 6: Annual rates of pertussis (per 100 000 population) by age group, <1 year vs. 1+ year, 1997-2011



Note: Rate of pertussis notified cases per 100 000 population calculated using mid-year population estimates.

Figure 7: Trends in cumulative pertussis rates (per 100 000 population) by age group and ethnicity, 2003 to 2011



Note: Rate of pertussis notified cases per 100 000 population (includes cases under investigation) calculated using mid-year population estimates



Figure 8: Monthly pertussis rates (cases per 100 000 population) by DHB, since January 2011





Figure 9: Monthly pertussis cases by DHB, since January 2011



**Note:** cases include confirmed, probable, and suspect only.

**Case classification for pertussis notification in New Zealand to 31 May 2012**

<b>Confirmed</b>	A clinically compatible illness that is laboratory confirmed by isolation of <i>Bordetella pertussis</i> from a pernasal swab, or epidemiologically linked to a confirmed case.
<b>Probable</b>	Cough lasting longer than two weeks and one or more of the following: <ul style="list-style-type: none"> <li>• Paroxysmal cough</li> <li>• Cough ending in vomiting or apnoea</li> <li>• Inspiratory whoop for which there is no other known cause.</li> </ul>
<b>Suspect</b>	In children under five years of age, any paroxysmal cough with whoop, vomiting or apnoea for which there is no other known cause.
<b>Other</b>	Status recorded as <i>under investigation</i> or suspect case.
<b>Notifications</b>	Include confirmed cases, probable, and other as specified above.

**Case classification for pertussis notification in New Zealand from 31 May 2012**

<b>Confirmed</b>	A clinically compatible illness that is laboratory confirmed by isolation of <i>B. pertussis</i> or detection of <i>B. pertussis</i> nucleic acid, preferably from a nasopharyngeal swab, or is epidemiologically linked to a confirmed case.
<b>Probable</b>	A clinically compatible illness with a high <i>B. pertussis</i> IgA test or a significant increase in antibody levels between paired sera at the same laboratory OR A cough lasting longer than two weeks and with one or more of the following, for which there is no other known cause: <ul style="list-style-type: none"> <li>• Paroxysmal cough</li> <li>• Cough ending in vomiting or apnoea</li> <li>• Inspiratory whoop</li> </ul>
<b>Suspect</b>	In children under five years of age any paroxysmal cough with whoop, vomiting or apnoea for which there is no other known cause.
<b>Under investigation</b>	A case that has been notified, but information is not yet available to classify it as suspect, probable or confirmed.
<b>Notifications</b>	Include confirmed cases, probable, suspect and under investigation as specified above.

This report will be available at: <http://www.surv.esr.cri.nz/surveillance/PertussisRpt.php>.