

# INFLUENZA WEEKLY UPDATE

2010/20: 17-23 May 2010

---

The national influenza surveillance system in New Zealand is an essential public health component for assessing and implementing strategies to control influenza. This report summarises the data collected from sentinel general practice (GP) surveillance and non-sentinel surveillance for week 20 (17-23 May 2010).

## SUMMARY OF THIS REPORT:

- Influenza-like illness (ILI) through sentinel surveillance was reported from 20 out of 21 District Health Boards (DHB) with a national consultation rate of 30.4 per 100 000 (131 ILI consultations). Sixty-one swabs were received from sentinel and non-sentinel surveillance. No influenza viruses were identified.
- Pandemic influenza A (H1N1) 09 is a notifiable disease in New Zealand. As of 23 May 2010, there have been 3306<sup>1</sup> cases recorded in EpiSurv with 35<sup>2</sup> deaths.

(Note: For further information please refer to the Ministry of Health's website:  
<http://www.moh.govt.nz/influenza-a-h1n1>)

In the past week, a total of 131 consultations for influenza-like illness were reported from 91 general practices in 20 out of 21 DHBs. This gives a weekly consultation rate of 30.4 per 100 000 patient population.


The graph below compares the consultation rates for influenza-like illness for each DHB over the past week. Counties Manukau DHB had the highest consultation rate (123.0 per 100 000, 92 cases), followed by Waikato (43.2 per 100 000, 8 cases) and Capital and Coast (22.7 per 100 000, 5 cases).

---

<sup>1</sup> Data source: EpiSurv as of 27 May 2010.

<sup>2</sup> Data source: Ministry of Health.

**Figure 1: Weekly consultation rates for influenza-like illness by DHB week ending 23 May 2010**




\*Caution is required in interpreting the ILI rate from Counties Manukau DHB because the contributing GP practices did not use the case definition of the sentinel influenza surveillance. The investigation to understand the specificity and sensitivity of their ILI case definition is ongoing.  
 [ ] No data for the week.

Sixty-one swabs were received by the virology laboratories from sentinel surveillance (14) and the non-sentinel surveillance (47). No influenza viruses were identified.

Figure 2 shows the weekly national consultation rates for 2008, 2009 seasons, and 2010 so far. The current rate of influenza-like illness is below the baseline.

**Figure 2: Weekly consultation rates for influenza-like illness in New Zealand, 2008, 2009 and 2010**



\* A weekly rate <50 ILI consultations per 100 000 patient population is considered baseline activity. A rate of 50–249 is considered indicative of normal seasonal influenza activity, and a rate of 250–399 indicative of higher than expected influenza activity. A rate >400 ILI consultations per 100 000 patient population indicates an epidemic level of influenza activity.

**Figure 3 illustrates consultation rates for influenza-like illness mapped by DHB for week 20, 2010**

