

# INFLUENZA WEEKLY UPDATE

2014/19: 5–11 May 2014

The national influenza surveillance system in New Zealand is an essential public health component for assessing and implementing strategies to control influenza. This report summarises the data collected from sentinel general practice (GP) surveillance and non-sentinel surveillance for week 19 (5–11 May 2014).

## Summary

- ILI through sentinel surveillance was reported from 17 out of 20 District Health Boards (DHB) with a national consultation rate of 16.4 per 100 000 (47 ILI consultations).
- A total of 227 swabs were received from sentinel (25) and non-sentinel (202) surveillance.
- 12 viruses were identified: A(H1N1)pdm09 (4), A(not sub-typed) (3), A(H3N2) (3), and B (lineage not determined) (2).

## INFLUENZA-LIKE ILLNESS SURVEILLANCE

In the past week, a total of 47 consultations for influenza-like illness (ILI) were reported from 55 general practices in 17 out of 20 DHBs. This gives a weekly consultation rate of 16.4 per 100 000 patient population. Figure 1 shows the weekly national consultation rate for 2014 in comparison to the average epidemic curve in 2000–2013 (excluding 2009). For more details on threshold definitions, see Appendix. The current rate of influenza-like illness is below the seasonal threshold.

**Figure 1. Weekly consultation rates for influenza-like illness in New Zealand in 2014 in comparison to the average epidemic curve in 2000–2013 (excluding 2009)**




Figure 2 shows the weekly national consultation rate for 2014 in comparison to the previous years 2010–2013.

**Figure 2. Weekly consultation rates for influenza-like illness in New Zealand, 2010–2014**






Figure 3 compares the consultation rates for influenza-like illness for each DHB over the past week. MidCentral DHB had the highest consultation rate (106.4 per 100 000, 3 cases) followed by and Hutt Valley (40.0 per 100 000, 6 cases).

**Figure 3. Weekly consultation rates for influenza-like illness by DHB week ending 11 May 2014**



\* No data for the week. [ ] Not participating in the influenza sentinel surveillance.  
 Based on the SHIVERS weekly report, the ILI incidence for Auckland and Counties Manukau DHBs for week 19 were 13.3 per 100 000 and 16.0 per 100 000 patient populations, respectively. For more details, please refer to the website: <http://www.esr.cri.nz/competencies/shivers/Pages/SHIVERSReports.aspx>

Figure 4. Consultation rates for influenza-like illness mapped by DHB for week 19, 2014



## VIROLOGICAL SURVEILLANCE

A total of 25 swabs were received from sentinel surveillance. Of these, two influenza viruses were identified as A(H1N1)pdm09 and A(H3N2) (1 each) from Canterbury DHB.


In addition, 202 swabs were received by virology laboratories from non-sentinel surveillance. Of these, 10 influenza viruses were identified: A (not sub-typed) (3), A(H1N1)pdm09 (3), A(H3N2) (2), and B (lineage not determined) (2). The distribution by DHB is shown in Table 1.

**Table 1. Influenza viruses from non-sentinel surveillance for week 19 by DHB**

Antigenic Strain	DHB				Total
	AK	CM	CC	CB	
A (not sub-typed)	2	0	1	0	3
A(H1N1)pdm09	0	1	0	2	3
A(H3N2)	1	0	0	1	2
B (lineage not determined)	0	2	0	0	2
<b>Total</b>	<b>3</b>	<b>3</b>	<b>1</b>	<b>3</b>	<b>10</b>

Figure 5 shows the cumulative total of influenza viruses confirmed (sentinel and non-sentinel surveillance) from week 1 to the end of week 19 (11 May 2014). A total of 203 influenza viruses were identified: A(H1N1)pdm09 (114) including seven A/California/7/2009 (H1N1)-like viruses, A (not sub-typed) (34), A(H3N2) (29) including 11 A/Victoria/361/2011 (H3N2)-like viruses, and influenza B (26) including six B/Wisconsin/1/2010- like viruses and one of B/Brisbane/60/2008-like.

**Figure 5. Cumulative laboratory-confirmed viruses by DHB from week 1 to week 19, 11 May 2014**



## APPENDIX

New Zealand's ILI data in recent years was reviewed and updated:

- The average epidemic curve (based on the 2000–2013 ILI data, excluding 2009) is the usual level of influenza activity that may occur during a typical year using the method described in “*Global epidemiological surveillance standards for influenza*” ([http://www.who.int/influenza/resources/documents/WHO\\_Epidemiological\\_Influenza\\_Surveillance\\_Standards\\_2014.pdf](http://www.who.int/influenza/resources/documents/WHO_Epidemiological_Influenza_Surveillance_Standards_2014.pdf)).
- The seasonal threshold is the level of influenza activity that signals the start and end of the annual influenza season and it was based on the 2000–2013 ILI data (excluding 2009) using the Moving Epidemic Method (*Vega et al. Influenza and other respiratory viruses 2013;7(4):546-558*). A weekly rate of 36 ILI consultations per 100 000 patient population is considered the seasonal threshold.
- Alert threshold (defined as 90% upper confidence interval of the mean) is a level above which, varying by time of year, influenza activity is higher than most years.
- A rate of 37–149 per 100 000 is used to describe normal seasonal influenza activity based on the 25th and 75th percentiles of the ILI data (2000–2013 excluding 2009). A rate of 150–399 is used to describe higher than expected influenza activity (i.e. 2009 pandemic). A rate of  $\geq 400$  is used to describe an epidemic level of influenza activity (i.e. 1996 experience).

Compiled by:

**Liza Lopez**

Health Intelligence Team  
ESR Kenepuru Science Centre  
PO Box 50 348, Porirua

T: 04 914 0647 F: 04 978 6690 E: [liza.lopez@esr.cri.nz](mailto:liza.lopez@esr.cri.nz)

**Dr. Sue Huang**

WHO National Influenza Centre  
ESR Wallaceville Science Centre  
PO Box 40158, Upper Hutt