



6SSN 2324-4097

# Community and Hospital Surveillance

## ILI, SARI, Influenza and Respiratory Pathogens

2017 Influenza Season, Week 35, ending 3 September 2017

### SUMMARY

- Influenza-like illness (ILI) consultation rates decreased compared to the previous week, and were below the seasonal threshold level.
- Influenza - associated severe acute respiratory illness (SARI) hospitalisation rates were low, also below the seasonal threshold level.
- Influenza A(H3N2) are the predominant viruses in New Zealand this year.
- Influenza B/Victoria lineage viruses also co-circulated with B/Yamagata lineage viruses with more B/Yamagata viruses this year.
- Rhinovirus and respiratory syncytial virus (RSV) are the most commonly detected non-influenza viruses this week, with numbers remaining high.

The surveillance for community-based influenza-like illness (ILI) and hospital-based severe acute respiratory illness (SARI) provides evidence to inform public health and clinical practice to reduce the impact of influenza virus infection and other important respiratory pathogens. This weekly report summarises data obtained from the ILI and SARI surveillance platforms. The report includes incidence, demographic characteristics, clinical outcomes and aetiologies for community ILI cases as well as hospital SARI cases including ICU admissions and deaths for the past week as well as the cumulative period since 2 January 2017.


Note: Data in this report are provisional and may change as more cases are assessed and information is updated. Data were extracted on 6 September 2017.

# INFLUENZA-LIKE ILLNESS and SEVERE ACUTE RESPIRATORY ILLNESS


## Influenza-like illness (ILI)

During week 35, ending 3 September 2017, 130 patients with influenza-like illness consulted sentinel general practices in 20 DHBs. The weekly ILI incidence was 31.6 per 100 000 patient population (Figures 1 and 2). Of the 72 tested ILI cases, 20 were positive for influenza viruses. This gives an ILI related influenza incidence (adjusted) of 8.8 per 100 000 patient population.

**Figure 1. Weekly resident ILI and influenza incidence since 2 January 2017**




**Figure 2. Comparison of 2017 rate with average seasonal rate, and historical thresholds**



The weekly consultation rates for influenza-like illness by different age groups and ethnicities are shown in Figures 3 and 4.

**Figure 3. Weekly ILI incidence by age group since 2 January 2017**



**Figure 4. Weekly ILI incidence by ethnicity since 2 January 2017**





Figure 5 compares the consultation rates for influenza-like illness for each DHB over the past week. Waitemata (316.4 per 100 000, 40 cases), Auckland (82.8 per 100 000, 47 cases), and Bay of Plenty (60.6 per 100 000, 9 cases) DHBs had the highest consultation rates.

**Figure 5. Rate of ILI consultations per 100 000 registered by DHB per week since 2 January 2017**



\*Results that have some uncertainty, with less than 5% of the DHB population covered (see Notes on Interpretation).  
Note: Outliers have been omitted from this graph.

ILI consultation rates for any particular DHB should be treated with caution as they may not be representative of the real situation for a particular community or setting, especially if the surveillance system has a small number of participating General Practices in the DHB, or the GP enrolled patient population is small, the calculated ILI rates are subject to greater fluctuation.

Since 2 January 2017, a total of 2495 ILI cases were identified. This gives a cumulative ILI incidence of 606.9 per 100 000 patient population (Table 1). Among the 1982 tested ILI cases, 888 (44.8%) were positive for influenza viruses. This gives an ILI related (adjusted) influenza incidence of 271.9 per 100 000 patient population.

**Table 1. Demographic characteristics of ILI and influenza cases, since 2 January 2017**

Characteristics	ILI & influenza cases among sentinel practices				
	ILI cases	Influenza cases	Prop Influenza positive <sup>1</sup> (%)	ILI incidence (per 100 000)	Influenza incidence <sup>2</sup> (per 100 000)
<b>Overall</b>	<b>2495</b>	<b>888</b>	<b>44.8 (100.0)</b>	<b>606.9</b>	<b>271.9</b>
<b>Age group (years)</b>					
<1	28	3	13.6 (0.3)	423.6	57.8
1–4	169	34	26.2 (3.8)	689.5	180.3
5–19	655	291	55.9 (32.8)	774.1	432.4
20–34	409	132	39.1 (14.9)	476.1	185.9
35–49	579	215	47.1 (24.2)	706.3	333.0
50–64	429	139	40.6 (15.7)	585.0	237.8
65–79	190	58	40.6 (6.5)	465.3	188.7
>80	36	16	53.3 (1.8)	269.5	143.7
Unknown	0	0	0.0		
<b>Ethnicity</b>					
Māori	242	82	43.4 (9.2)	425.2	184.5
Pacific peoples	111	42	48.3 (4.7)	371.2	179.2
Asian	241	119	54.6 (13.4)	804.4	439.1
European and Other	1899	645	43.4 (72.6)	646.7	280.5
Unknown	2	0	0.0	0.0	
<b>Sex</b>					
Female	1419	494	44.1 (55.6)	664.9	293.0
Male	1072	394	45.8 (44.4)	542.2	248.1
Unknown	4	0	0.0		


<sup>1</sup>Proportion of cases tested which were positive for influenza viruses

<sup>2</sup>Adjusted to positivity of tested cases

## Severe acute respiratory illness (SARI)

There were 2832 acute admissions to ADHB and CMDHB hospitals during week 35, ending 3 September 2017. A total of 127 patients with acute respiratory illness were assessed in these hospitals. Of these, 83 (65.4%) patients met the SARI case definition. Five cases were admitted to ICU and no SARI related deaths were reported this week. Of the 52 tested residents with SARI, six were positive for influenza viruses, giving a SARI related influenza incidence of 0.8 per 100 000 population this week.

**Figure 6. Weekly resident SARI and influenza incidence since 2 May 2016 and previous seasons SARI incidence**



Since 1 May 2017, a total of 1450 SARI cases were identified. This gives a SARI proportion of 28.4 per 1000 acute hospitalisations (Table 2). Ninety-eight SARI cases have been admitted to ICU and 22 SARI related deaths were reported during this period.

Of the 1450 SARI cases, 1123 were ADHB and CMDHB residents, giving a SARI incidence of 124.0 per 100 000 population (Table 2). Among the 1020 tested SARI cases who were ADHB and CMDHB residents, 325 (31.9%) had positive influenza virus results. This gives a SARI (adjusted) related influenza incidence of 39.5 per 100 000 population.

**Table 2. Demographic characteristics of SARI cases and related influenza cases, since 1 May 2017**

Characteristics	Admissions	Assessed	SARI & influenza cases among all hospital patients			SARI & influenza cases among ADHB & CMDHB residents			
			SARI Cases (%)	Cases per 1000 hospitalisations	Influenza positive <sup>1</sup> (%)	SARI cases	SARI incidence (per 100 000)	Influenza Cases	Influenza incidence (per 100 000)
<b>Overall</b>	<b>51075</b>	<b>2680</b>	<b>1450 (54.1)</b>	<b>28.4</b>	<b>346 (31.8)</b>	<b>1123</b>	<b>124.0</b>	<b>325</b>	<b>39.5</b>
<b>Age group (years)</b>									
<1	1976		252	127.5	18 (8.2)	235	1740.0	17	143.6
1–4	3566		171	48.0	30 (21.4)	157	296.9	28	64.9
5–19	6039		77	12.8	18 (27.7)	63	32.7	12	7.3
20–34	9561		70	7.3	28 (41.8)	67	32.1	27	13.6
35–49	7438		84	11.3	32 (40.5)	79	41.4	31	17.3
50–64	8680		177	20.4	79 (47.3)	172	114.3	77	54.3
65–79	8287		215	25.9	75 (36.9)	204	279.1	69	100.3
>80	5528		150	27.1	66 (45.8)	146	623.1	64	284.9
Unknown	0		254			0		0	
<b>Ethnicity</b>									
Māori	6920		242	35.0	46 (21.3)	218	219.2	43	48.6
Pacific peoples	10957		423	38.6	114 (29.7)	410	297.1	107	85.7
Asian	8493		101	11.9	33 (37.1)	97	46.1	32	16.8
European and Other	24346		430	17.7	153 (38.7)	398	99.1	143	38.6
Unknown	359		254	707.5		0		0	
<b>Hospitals</b>									
ADHB	29830	1007	706 (70.1)	23.7	173 (37.9)	447	102.4	155	39.4
CMDHB	21245	1673	744 (44.5)	35.0	173 (27.4)	676	144.0	170	39.7
<b>Sex</b>									
Female	27049		612	22.6	181 (32.7)	570	122.5	171	40.5
Male	24023		580	24.1	164 (31.1)	550	124.9	153	38.3
Unknown	3		258			3		1	

<sup>1</sup>Proportion of cases tested which were positive for influenza viruses

Note. A specimen may be positive for more than one virus; a patient may have more than one specimen tested.

## RESPIRATORY PATHOGEN SURVEILLANCE

### Influenza virus

During week 35, 72 ILI specimens were tested; 20 were positive for influenza viruses. In addition, 65 SARI specimens were tested; six were positive for influenza viruses.

Since 1 May 2017, 1967 ILI specimens were tested, 887 (45.1%) were positive for influenza with the following viruses. In addition, 1228 SARI specimens were tested, 383 (31.2%) were positive for influenza viruses (see Table 3).

**Table 3. Influenza viruses among ILI and SARI cases since 1 May 2017**

Influenza viruses	ILI	SARI	SARI and non-SARI	
	Cases (%)	Cases (%)	ICU (%)	Deaths (%)
No. of specimens tested	1967	1228	218	25
No. of positive specimens (%) <sup>1</sup>	887 (45.1)	383 (31.2)	24 (11.0)	12 (48.0)
<b>Influenza A</b>	<b>517</b>	<b>284</b>	<b>15</b>	<b>7</b>
A (not subtyped)	12	109	5	1
A(H1N1)pdm09	67	33	3	0
A(H1N1)pdm09 by PCR	49	29	2	0
A/Michigan/45/2015 (H1N1)pdm09 - like	14	4	1	0
A/California/7/2009 (H1N1)pdm09 - like	4	0	0	0
A(H3N2)	438	142	7	6
A(H3N2) by PCR	389	140	7	6
A/Hong Kong/4801/2014 (H3N2) - like	49	2	0	0
<b>Influenza B</b>	<b>370</b>	<b>99</b>	<b>9</b>	<b>5</b>
B (lineage not determined)	17	53	7	4
B/Yamagata lineage	338	45	2	1
B/Yamagata lineage by PCR	174	22	2	1
B/Phuket/3073/2013 - like	164	23	0	0
B/Victoria lineage	15	1	0	0
B/Victoria lineage by PCR	7	1	0	0
B/Brisbane/60/2008 - like	8	0	0	0
<b>Influenza and non-influenza co-detection (% +ve)</b>	<b>50 (5.6)</b>	<b>24 (6.3)</b>	<b>3 (12.5)</b>	<b>1 (8.3)</b>

<sup>1</sup>Number of specimens positive for at least one of the listed viruses

Note. A specimen may be positive for more than one virus; a patient may have more than one specimen tested.

The recommended influenza vaccine formulation for trivalent vaccine for New Zealand in 2017 is:

- A(H1N1) an A/Michigan/45/2015 (H1N1)pdm09-like virus
- A(H3N2) an A/Hong Kong/4801/2014 (H3N2)-like virus
- B a B/Brisbane/60/2008-like virus (belonging to B/Victoria lineage)

Quadrivalent vaccines contain the above three viruses plus one more vaccine component: B/Phuket/3073/2013-like virus (belonging to B/Yamagata lineage)

Note: Antigenic characterization of the current A(H3N2) viruses have been technically challenging because many viruses had low or undetectable haemagglutination activity. This phenomenon has been well recognized globally and documented in WHO's Weekly Epidemiological Record:

<http://apps.who.int/iris/bitstream/10665/254756/1/WER9211.pdf?ua=1>



## Non-influenza respiratory pathogens


Since 1 May 2017, 1918 ILI specimens were tested for non-influenza viruses, 399 (20.8%) were positive with the following viruses. One thousand and eight SARI specimens were tested for non-influenza viruses, 341 (33.8%) were positive with the following viruses (see Table 4).

**Table 4. Non-influenza viruses among ILI and SARI cases since 1 May 2017<sup>1</sup>**


<i>Non-influenza respiratory viruses</i>	ILI	SARI	SARI and non-SARI	
	Cases (%)	Cases (%)	ICU (%)	Deaths (%)
No. of specimens tested	1918	1008	164	24
No. of positive specimens (%) <sup>1</sup>	399 (20.8)	341 (33.8)	104 (63.4)	1 (4.2)
Respiratory syncytial virus (RSV)	108	183	53	1
Parainfluenza 1 (PIV1)	3	1	0	0
Parainfluenza 2 (PIV2)	23	13	4	0
Parainfluenza 3 (PIV3)	49	32	8	0
Rhinovirus (RV)	132	112	48	0
Adenovirus (AdV)	49	38	11	0
Human metapneumovirus (hMPV)	32	20	3	0
Enterovirus	26	19	8	0
Single virus detection (% of positives)	379 (95.0)	274 (80.4)	77 (74.0)	0 (-)
Multiple virus detection (% of positives)	20 (5.0)	67 (19.6)	27 (26.0)	0 (-)

<sup>1</sup>Number of specimens positive for at least one of the listed viruses; note a specimen may be positive for more than one virus


**Figure 7. Temporal distribution of the number and proportion of influenza viruses from ILI specimens by type and week<sup>1</sup>**




**Figure 8. Temporal distribution of the number and proportion of influenza viruses from SARI specimens by type and week<sup>1</sup>**



**Figure 9. Temporal distribution of the number and proportion of non-influenza viruses from ILI specimens by type and week<sup>1</sup>**



**Figure 10. Temporal distribution of the number and proportion of non-influenza viruses from SARI specimens by type and week<sup>1</sup>**



<sup>1</sup>Figures for recent weeks will be underestimates due to time lag in receiving laboratory test results.



## APPENDIX

Table 5. Influenza-like illness count by DHB by week 18–35, 2017

DHB	Week																	
	18	19	20	21	22	23	24	25	26	27	28	29	30	31	32	33	34	35
Auckland	15	14	15	18	30	25	45	71	58	83	78	62	82	69	58	60	64	47
Bay of Plenty	0	0	0	0	2	1	1	6	7	11	4	4	5	6	7	7	11	9
Canterbury	2	5	10	3	3	1	11	16	9	12	9	20	18	15	22	18	20	5
Capital and Coast	2	2	3	4	11	9	13	6	8	25	18	22	13	7	16	23	7	5
Counties Manukau	1	2	0	2	1	0	3	1	6	3	6	1	2	3	0	1	0	0
Hawke's Bay	0	0	1	3	1	1	3	0	4	4	8	2	3	8	7	3	1	2
Hutt Valley	4	0	0	0	0	1	1	1	0	0	0	0	0	1	0	0	0	0
Lakes	0	0	0	1	0	0	0	0	0	0	3	0	0	0	0	0	0	0
MidCentral	0	0	0	0	0	0	0	1	1	1	0	0	1	1	1	0	1	1
Nelson Marlborough	0	3	2	0	1	0	4	3	2	2	4	9	3	4	7	3	2	0
Northland	2	1	4	2	0	0	1	8	4	4	2	2	0	0	1	0	0	2
South Canterbury	1	3	1	2	0	0	2	1	1	1	0	5	10	8	8	16	23	6
Southern	1	2	0	5	4	5	1	2	4	4	7	11	4	10	11	4	6	5
Tairāwhiti	0	0	1	2	4	2	2	7	3	2	2	4	2	2	2	2	1	0
Taranaki	1	0	0	1	0	0	1	3	1	2	1	1	0	0	2	1	0	1
Waikato	0	5	3	3	4	6	4	8	4	13	9	6	5	3	7	4	1	7
Wairarapa	0	0	0	0	0	0	0	0	5	2	1	1	0	0	0	0	0	0
Waitemata	1	6	2	8	22	16	24	44	41	37	30	44	50	52	37	39	24	40
West Coast	1	1	0	0	0	2	0	1	3	2	0	0	0	1	2	0	0	0
Whanganui	0	0	1	4	0	1	2	0	5	5	5	0	6	4	1	1	1	0
<b>New Zealand</b>	<b>31</b>	<b>44</b>	<b>43</b>	<b>58</b>	<b>83</b>	<b>70</b>	<b>118</b>	<b>179</b>	<b>166</b>	<b>213</b>	<b>187</b>	<b>194</b>	<b>204</b>	<b>194</b>	<b>189</b>	<b>182</b>	<b>162</b>	<b>130</b>

**Table 6. Influenza-like illness rate by DHB by week 18–35, 2017**

DHB	Rate per 100 000																	
	18	19	20	21	22	23	24	25	26	27	28	29	30	31	32	33	34	35
Auckland	26.4	24.7	26.4	31.7	52.8	44.0	79.2	125.0	102.1	146.2	137.4	109.2	144.4	121.5	102.1	105.7	112.7	82.8
Bay of Plenty	0.0	0.0	0.0	0.0	13.5	6.7	6.7	40.4	47.1	74.1	26.9	26.9	33.7	40.4	47.1	47.1	74.1	60.6
Canterbury	2.8	7.1	14.2	4.3	4.3	1.4	15.6	22.8	12.8	17.1	12.8	28.4	25.6	21.3	31.3	25.6	28.4	7.1
Capital and Coast	7.8	7.8	11.7	15.6	42.8	35.0	50.6	23.4	31.2	97.3	70.1	85.7	50.6	27.3	62.3	89.6	27.3	19.5
Counties Manukau	3.0	6.0	0.0	6.0	3.0	0.0	9.0	3.0	18.0	9.0	18.0	3.0	6.0	9.0	0.0	3.0	0.0	0.0
Hawke's Bay	0.0	0.0	5.2	15.7	5.2	5.2	15.7	0.0	20.9	20.9	41.7	10.4	15.7	41.7	36.5	15.7	5.2	10.4
Hutt Valley	15.2	0.0	0.0	0.0	0.0	3.8	3.8	3.8	0.0	0.0	0.0	0.0	0.0	3.8	0.0	0.0	0.0	0.0
Lakes*	0.0	0.0	0.0	22.3	0.0	0.0	0.0	0.0	0.0	0.0	66.8	0.0	0.0	0.0	0.0	0.0	0.0	0.0
MidCentral*	0.0	0.0	0.0	0.0	0.0	0.0	0.0	35.5	35.5	35.5	0.0	0.0	35.5	35.5	35.5	0.0	35.5	35.5
Nelson Marlborough	0.0	29.2	19.5	0.0	9.7	0.0	39.0	29.2	19.5	19.5	39.0	87.7	29.2	39.0	68.2	29.2	19.5	0.0
Northland	15.7	7.9	31.5	15.7	0.0	0.0	7.9	62.9	31.5	31.5	15.7	15.7	0.0	0.0	7.9	0.0	0.0	15.7
South Canterbury	9.3	27.8	9.3	18.6	0.0	0.0	18.6	9.3	9.3	9.3	0.0	46.4	92.8	74.2	74.2	148.5	213.4	55.7
Southern	2.2	4.4	0.0	10.9	8.7	10.9	2.2	4.4	8.7	8.7	15.3	24.0	8.7	21.8	24.0	8.7	13.1	10.9
Tairāwhiti	0.0	0.0	14.4	28.9	57.7	28.9	28.9	101.0	43.3	28.9	28.9	57.7	28.9	28.9	28.9	28.9	14.4	0.0
Taranaki	4.4	0.0	0.0	4.4	0.0	0.0	4.4	13.1	4.4	8.7	4.4	4.4	0.0	0.0	8.7	4.4	0.0	4.4
Waikato*	0.0	32.5	19.5	19.5	26.0	39.0	26.0	52.1	26.0	84.6	58.6	39.0	32.5	19.5	45.5	26.0	6.5	45.5
Wairarapa	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	200.0	80.0	40.0	40.0	0.0	0.0	0.0	0.0	0.0	0.0
Waitematā*	7.9	47.5	15.8	63.3	174.0	126.6	189.8	348.0	324.3	292.7	237.3	348.0	395.5	411.3	292.7	308.5	189.8	316.4
West Coast	6.3	6.3	0.0	0.0	0.0	12.7	0.0	6.3	19.0	12.7	0.0	0.0	0.0	6.3	12.7	0.0	0.0	0.0
Whanganui*	0.0	0.0	60.2	241.0	0.0	60.2	120.5	0.0	301.2	301.2	301.2	0.0	361.4	241.0	60.2	60.2	60.2	0.0
<b>New Zealand</b>	<b>7.5</b>	<b>10.7</b>	<b>10.5</b>	<b>14.1</b>	<b>20.2</b>	<b>17.0</b>	<b>28.7</b>	<b>43.5</b>	<b>40.4</b>	<b>51.8</b>	<b>45.5</b>	<b>47.2</b>	<b>49.6</b>	<b>47.2</b>	<b>46.0</b>	<b>44.3</b>	<b>39.4</b>	<b>31.6</b>

\*Results that have some uncertainty, with less than 5% of the DHB population covered (see Notes on Interpretation).



Recent global experience with pandemic influenza A(H1N1)pdm09 highlights the importance of monitoring severe and mild respiratory disease to support pandemic preparedness as well as seasonal influenza prevention and control. Two active, prospective, population-based surveillance systems were used to monitor influenza and other respiratory pathogens: 1) among those registered patients seeking consultations with influenza-like illness (ILI) at sentinel general practices nation-wide; 2) among those hospitalized patients with severe acute respiratory illness (SARI) in Auckland and Counties Manukau District Health Boards (ADHB and CMDHB).

The aims of ILI and SARI surveillance are: 1) to measure the burden of severe and moderate disease caused by influenza and other respiratory pathogens; 2) to monitor trends in severe and moderate disease caused by influenza and other respiratory pathogens; 3) to identify high risk groups that should be prioritized for prevention and treatment; 4) to monitor antigenic, genetic and antiviral characteristics of influenza viruses associated with severe and mild disease. 5) to provide a study base to estimate the effectiveness of influenza vaccine.

## ACKNOWLEDGEMENT

We acknowledge the support of the New Zealand Ministry of Health. SARI surveillance was established and funded by the US CDC, and continues to operate through funding from the New Zealand Ministry of Health.

## DESCRIPTION OF ILI ACTIVITY THRESHOLDS

The values for the different intensity levels for 2017 are listed in the table below. This is based on New Zealand’s consultation rates from 2000–2015 (excluding the pandemic year, 2009) and WHO’s interim guidance severity assessment

Below seasonal level (baseline, per 100,000)	Seasonal level (per 100,000)			Above seasonal level (per 100,000)
	low	moderate	high	
<35.1	35.1-82.5	82.5-168.9	168.9-231.8	>231.8

- The baseline threshold indicates the level of influenza activity that signals the start and end of the annual influenza season and it is based on the Moving Epidemic Method (MEM) (*Vega et al. Influenza and other respiratory viruses 2013;7(4):546-558*).
- Seasonal levels (low, moderate and high) are estimated as the upper limits of the 40%, 90% and 97.5% one-sided confidence intervals of the geometric mean of 30 highest epidemic weekly rates using the MEM method. As many other countries use this method, it allows the NZ data to be interpreted not just at the country level but also comparable with other countries.
- The average seasonal curve indicates the usual seasonal activity that may occur during a typical year using the method described in “*Global epidemiological surveillance standards for influenza*” ([http://www.who.int/influenza/resources/documents/WHO\\_Epidemiological\\_Influenza\\_Surveillance\\_Standards\\_2014.pdf](http://www.who.int/influenza/resources/documents/WHO_Epidemiological_Influenza_Surveillance_Standards_2014.pdf)).

## NOTES ON INTERPRETATION

- SARI case definition: “An acute respiratory illness with a history of fever or measured fever of  $\geq 38^{\circ}\text{C}$ , AND cough, AND onset within the past 10 days, AND requiring inpatient hospitalisation (defined as a patient who is admitted under a medical team and to a hospital ward or assessment unit)”. A non-SARI case is a hospitalised respiratory patient who does not meet the SARI case definition.
- ILI case definition: “An acute respiratory illness with a history of fever or measured fever of  $\geq 38^{\circ}\text{C}$ , AND cough, AND onset within the past 10 days, AND requiring GP consultation”.
- ILI sentinel general practices: a total of 74 sentinel general practices have agreed to participate in community ILI surveillance. These practices have ~400 000 registered patients, covering roughly 9% of the NZ population.
- SARI sentinel hospitals serving a population of 906 000 people: Auckland City Hospital and the associated Starship Children’s Hospital (ADHB), and Middlemore Hospital and the associated Kidz First Children’s Hospital (CMDHB).
- The real-time PCR assay for influenza virus uses CDC’s protocol ([http://www.accessdata.fda.gov/cdrh\\_docs/pdf8/k080570.pdf](http://www.accessdata.fda.gov/cdrh_docs/pdf8/k080570.pdf));
- The real-time PCR assay for non-influenza respiratory viruses (respiratory syncytial virus, parainfluenza virus types 1-3, human metapneumovirus, rhinovirus and adenovirus) uses CDC’s protocol. Note: The rhinovirus PCR detects mostly rhinovirus with slight cross-reactivity against enterovirus.
- The surveillance week is Monday to Sunday inclusive, and data are extracted on the subsequent Tuesday. Results from previous weeks will be revised as data are updated (laboratory test results in particular may be delayed).
- ILI consultation rates for any particular DHB should be treated with caution. If the surveillance system has a small number of participating General Practices in the DHB, or the GP enrolled patient population is small, the calculated ILI rates are subject to greater fluctuation.

This weekly report is compiled by ESR. For more information please contact:

Tim Wood: T:+64 4 529 0611; E: [Tim.Wood@esr.cri.nz](mailto:Tim.Wood@esr.cri.nz)

Liza Lopez: T:+64 4 914 0647; E: [Liza.Lopez@esr.cri.nz](mailto:Liza.Lopez@esr.cri.nz)

Sue Huang: T:+64 4 529 0606; E: [Sue.Huang@esr.cri.nz](mailto:Sue.Huang@esr.cri.nz)