



ISSN 2324-4097

# Community and Hospital Surveillance

## ILI, SARI, Influenza and Respiratory Pathogens

2017 Influenza Season, Week 37, ending 17 September 2017

### SUMMARY

- Influenza-like illness (ILI) consultation rates decreased compared to the previous week, and were below the seasonal threshold level.
- Influenza - associated severe acute respiratory illness (SARI) hospitalisation rates were similar to last week, at a low level and below the seasonal threshold level.
- Influenza A(H3N2) are the predominant viruses in New Zealand this year.
- Influenza B/Victoria lineage viruses also co-circulated with B/Yamagata lineage viruses with more B/Yamagata viruses this year.
- Rhinovirus and respiratory syncytial virus (RSV) are the most commonly detected non-influenza viruses this week, with numbers remaining high.

The surveillance for community-based influenza-like illness (ILI) and hospital-based severe acute respiratory illness (SARI) provides evidence to inform public health and clinical practice to reduce the impact of influenza virus infection and other important respiratory pathogens. This weekly report summarises data obtained from the ILI and SARI surveillance platforms. The report includes incidence, demographic characteristics, clinical outcomes and aetiologies for community ILI cases as well as hospital SARI cases including ICU admissions and deaths for the past week as well as the cumulative period since 2 January 2017.


Note: Data in this report are provisional and may change as more cases are assessed and information is updated. Data were extracted on 20 September 2017.

# INFLUENZA-LIKE ILLNESS and SEVERE ACUTE RESPIRATORY ILLNESS


## Influenza-like illness (ILI)

During week 37, ending 17 September 2017, 106 patients with influenza-like illness consulted sentinel general practices in 20 DHBs. The weekly ILI incidence was 25.8 per 100 000 patient population (Figures 1 and 2). Of the 58 tested ILI cases, 12 were positive for influenza viruses. This gives an ILI related influenza incidence (adjusted) of 5.3 per 100 000 patient population.

**Figure 1. Weekly resident ILI and influenza incidence since 2 January 2017**




**Figure 2. Comparison of 2017 rate with average seasonal rate, and historical thresholds**



The weekly consultation rates for influenza-like illness by different age groups and ethnicities are shown in Figures 3 and 4.

**Figure 3. Weekly ILI incidence by age group since 2 January 2017**



**Figure 4. Weekly ILI incidence by ethnicity since 2 January 2017**





Figure 5 compares the consultation rates for influenza-like illness for each DHB over the past week. Waitemata (205.6 per 100 000, 26 cases), Auckland (86.3 per 100 000, 49 cases), and South Canterbury (83.5 per 100 000, 9 cases) DHBs had the highest consultation rates.

**Figure 5. Rate of ILI consultations per 100 000 registered by DHB per week since 2 January 2017**



\*Results that have some uncertainty, with less than 5% of the DHB population covered (see Notes on Interpretation).  
 Note: Outliers have been omitted from this graph.

ILI consultation rates for any particular DHB should be treated with caution as they may not be representative of the real situation for a particular community or setting, especially if the surveillance system has a small number of participating General Practices in the DHB, or the GP enrolled patient population is small, the calculated ILI rates are subject to greater fluctuation.

Since 2 January 2017, a total of 2712 ILI cases were identified. This gives a cumulative ILI incidence of 659.6 per 100 000 patient population (Table 1). Among the 2171 tested ILI cases, 931 (42.9%) were positive for influenza viruses. This gives an ILI related (adjusted) influenza incidence of 282.9 per 100 000 patient population.

**Table 1. Demographic characteristics of ILI and influenza cases, since 2 January 2017**

Characteristics	ILI & influenza cases among sentinel practices				
	ILI cases	Influenza cases	Prop Influenza positive <sup>1</sup> (%)	ILI incidence (per 100 000)	Influenza incidence <sup>2</sup> (per 100 000)
<b>Overall</b>	<b>2712</b>	<b>931</b>	<b>42.9 (100.0)</b>	<b>659.6</b>	<b>282.9</b>
<b>Age group (years)</b>					
<1	37	3	10.7 (0.3)	559.8	60.0
1–4	181	36	25.9 (3.9)	738.5	191.3
5–19	709	305	53.0 (32.8)	838.0	444.5
20–34	440	135	37.1 (14.5)	512.2	189.9
35–49	633	230	46.1 (24.7)	772.2	355.9
50–64	465	146	39.4 (15.7)	634.1	249.5
65–79	208	59	36.6 (6.3)	509.3	186.7
>80	39	17	50.0 (1.8)	292.0	146.0
Unknown	0	0	0.0		
<b>Ethnicity</b>					
Māori	256	84	42.2 (9.0)	449.8	189.9
Pacific peoples	117	43	47.8 (4.6)	391.3	187.0
Asian	262	123	51.9 (13.2)	874.4	453.8
European and Other	2075	681	41.4 (73.1)	706.6	292.7
Unknown	2	0	0.0	0.0	
<b>Sex</b>					
Female	1547	520	42.2 (55.9)	724.8	306.2
Male	1161	411	43.7 (44.1)	587.2	256.8
Unknown	4	0	0.0		


<sup>1</sup>Proportion of cases tested which were positive for influenza viruses

<sup>2</sup>Adjusted to positivity of tested cases

## Severe acute respiratory illness (SARI)

There were 2701 acute admissions to ADHB and CMDHB hospitals during week 37, ending 17 September 2017. A total of 115 patients with acute respiratory illness were assessed in these hospitals. Of these, 73 (63.5%) patients met the SARI case definition. Four cases were admitted to ICU and no SARI related deaths were reported this week. Of the 50 tested residents with SARI, eight were positive for influenza viruses, giving a SARI related influenza incidence of 1.1 per 100 000 population this week.

**Figure 6. Weekly resident SARI and influenza incidence since 2 May 2016 and previous seasons SARI incidence**



Since 1 May 2017, a total of 1616 SARI cases were identified. This gives a SARI proportion of 28.5 per 1000 acute hospitalisations (Table 2). One hundred and nine SARI cases have been admitted to ICU and 26 SARI related deaths were reported during this period.

Of the 1616 SARI cases, 1260 were ADHB and CMDHB residents, giving a SARI incidence of 139.1 per 100 000 population (Table 2). Among the 1144 tested SARI cases who were ADHB and CMDHB residents, 342 (29.9%) had positive influenza virus results. This gives a SARI (adjusted) related influenza incidence of 41.6 per 100 000 population.

**Table 2. Demographic characteristics of SARI cases and related influenza cases, since 1 May 2017**

Characteristics	Admissions	Assessed	SARI & influenza cases among all hospital patients			SARI & influenza cases among ADHB & CMDHB residents			
			SARI Cases (%)	Cases per 1000 hospitalisations	Influenza positive <sup>1</sup> (%)	SARI cases	SARI incidence (per 100 000)	Influenza Cases	Influenza incidence (per 100 000)
<b>Overall</b>	<b>56625</b>	<b>2947</b>	<b>1616 (54.8)</b>	<b>28.5</b>	<b>365 (30.0)</b>	<b>1260</b>	<b>139.1</b>	<b>342</b>	<b>41.6</b>
<b>Age group (years)</b>									
<1	2242		295	131.6	21 (8.2)	277	2050.9	20	168.8
1–4	3909		195	49.9	32 (20.4)	177	334.7	30	70.7
5–19	6673		87	13.0	21 (29.2)	72	37.4	15	9.2
20–34	10642		80	7.5	29 (38.2)	77	36.9	28	14.2
35–49	8245		89	10.8	32 (38.1)	82	42.9	30	16.7
50–64	9617		188	19.5	81 (45.8)	183	121.6	79	55.8
65–79	9216		242	26.3	81 (35.1)	230	314.7	74	105.9
>80	6081		165	27.1	68 (42.8)	161	687.2	66	292.6
Unknown	0		274			0		0	
<b>Ethnicity</b>									
Māori	7731		278	36.0	51 (20.5)	250	251.3	46	51.6
Pacific peoples	12089		467	38.6	121 (28.5)	453	328.3	114	91.1
Asian	9467		112	11.8	36 (37.1)	108	51.3	35	18.7
European and Other	26946		483	17.9	157 (35.5)	448	111.6	147	39.7
Unknown	392		276	704.1		1		0	
<b>Hospitals</b>									
ADHB	33070	1095	781 (71.3)	23.6	178 (35.5)	500	114.6	160	41.1
CMDHB	23555	1852	835 (45.1)	35.4	187 (26.2)	760	161.9	182	42.2
<b>Sex</b>									
Female	30024		688	22.9	193 (30.9)	642	138.0	183	42.9
Male	26598		651	24.5	170 (29.1)	616	139.9	158	40.0
Unknown	3		277			2		1	

<sup>1</sup>Proportion of cases tested which were positive for influenza viruses

Note. A specimen may be positive for more than one virus; a patient may have more than one specimen tested.

## RESPIRATORY PATHOGEN SURVEILLANCE

### Influenza virus

During week 37, 58 ILI specimens were tested; 12 were positive for influenza viruses. In addition, 54 SARI specimens were tested; eight were positive for influenza viruses.

Since 1 May 2017, 2156 ILI specimens were tested, 930 (43.1%) were positive for influenza with the following viruses. In addition, 1373 SARI specimens were tested, 403 (29.4%) were positive for influenza viruses (see Table 3).

**Table 3. Influenza viruses among ILI and SARI cases since 1 May 2017**

Influenza viruses	ILI	SARI	SARI and non-SARI	
	Cases (%)	Cases (%)	ICU (%)	Deaths (%)
No. of specimens tested	2156	1373	246	28
No. of positive specimens (%) <sup>1</sup>	930 (43.1)	403 (29.4)	24 (9.8)	13 (46.4)
<b>Influenza A</b>	<b>536</b>	<b>295</b>	<b>15</b>	<b>8</b>
A (not subtyped)	14	114	5	1
A(H1N1)pdm09	67	36	3	0
A(H1N1)pdm09 by PCR	47	32	2	0
A/Michigan/45/2015 (H1N1)pdm09 - like	16	4	1	0
A/California/7/2009 (H1N1)pdm09 - like	4	0	0	0
A(H3N2)	455	145	7	7
A(H3N2) by PCR	404	141	7	7
A/Hong Kong/4801/2014 (H3N2) - like	51	4	0	0
<b>Influenza B</b>	<b>394</b>	<b>108</b>	<b>9</b>	<b>5</b>
B (lineage not determined)	27	60	7	4
B/Yamagata lineage	352	47	2	1
B/Yamagata lineage by PCR	166	22	2	1
B/Phuket/3073/2013 - like	186	25	0	0
B/Victoria lineage	15	1	0	0
B/Victoria lineage by PCR	7	1	0	0
B/Brisbane/60/2008 - like	8	0	0	0
<b>Influenza and non-influenza co-detection (% +ve)</b>	<b>54 (5.8)</b>	<b>26 (6.5)</b>	<b>3 (12.5)</b>	<b>1 (7.7)</b>

<sup>1</sup>Number of specimens positive for at least one of the listed viruses

Note. A specimen may be positive for more than one virus; a patient may have more than one specimen tested.

The recommended influenza vaccine formulation for trivalent vaccine for New Zealand in 2017 is:

- A(H1N1) an A/Michigan/45/2015 (H1N1)pdm09-like virus
- A(H3N2) an A/Hong Kong/4801/2014 (H3N2)-like virus
- B a B/Brisbane/60/2008-like virus (belonging to B/Victoria lineage)

Quadrivalent vaccines contain the above three viruses plus one more vaccine component: B/Phuket/3073/2013-like virus (belonging to B/Yamagata lineage)

Note: Antigenic characterization of the current A(H3N2) viruses have been technically challenging because many viruses had low or undetectable haemagglutination activity. This phenomenon has been well recognized globally and documented in WHO's Weekly Epidemiological Record: <http://apps.who.int/iris/bitstream/10665/254756/1/WER9211.pdf?ua=1>



## Non-influenza respiratory pathogens


Since 1 May 2017, 2121 ILI specimens were tested for non-influenza viruses, 476 (22.4%) were positive with the following viruses. One thousand and one hundred sixty-five SARI specimens were tested for non-influenza viruses, 410 (35.2%) were positive with the following viruses (see Table 4).

**Table 4. Non-influenza viruses among ILI and SARI cases since 1 May 2017<sup>1</sup>**


<i>Non-influenza respiratory viruses</i>	ILI	SARI	SARI and non-SARI	
	Cases (%)	Cases (%)	ICU (%)	Deaths (%)
No. of specimens tested	2121	1165	187	26
No. of positive specimens (%) <sup>1</sup>	476 (22.4)	410 (35.2)	121 (64.7)	1 (3.8)
Respiratory syncytial virus (RSV)	125	213	63	1
Parainfluenza 1 (PIV1)	5	1	0	0
Parainfluenza 2 (PIV2)	24	15	4	0
Parainfluenza 3 (PIV3)	58	43	8	0
Rhinovirus (RV)	157	134	54	0
Adenovirus (AdV)	59	42	11	0
Human metapneumovirus (hMPV)	44	29	4	0
Enterovirus	35	26	11	0
Single virus detection (% of positives)	449 (94.3)	328 (80.0)	91 (75.2)	0 (-)
Multiple virus detection (% of positives)	27 (5.7)	82 (20.0)	30 (24.8)	0 (-)

<sup>1</sup>Number of specimens positive for at least one of the listed viruses; note a specimen may be positive for more than one virus


**Figure 7. Temporal distribution of the number and proportion of influenza viruses from ILI specimens by type and week<sup>1</sup>**




**Figure 8. Temporal distribution of the number and proportion of influenza viruses from SARI specimens by type and week<sup>1</sup>**



**Figure 9. Temporal distribution of the number and proportion of non-influenza viruses from ILI specimens by type and week<sup>1</sup>**



**Figure 10. Temporal distribution of the number and proportion of non-influenza viruses from SARI specimens by type and week<sup>1</sup>**



<sup>1</sup>Figures for recent weeks will be underestimates due to time lag in receiving laboratory test results.

## APPENDIX

Table 5. Influenza-like illness count by DHB by week 18–37, 2017

DHB	Week																			
	18	19	20	21	22	23	24	25	26	27	28	29	30	31	32	33	34	35	36	37
Auckland	15	14	15	18	30	25	45	71	58	83	78	62	82	69	58	60	64	47	38	49
Bay of Plenty	0	0	0	0	2	1	1	6	7	11	4	4	5	6	7	7	11	9	2	2
Canterbury	2	5	10	3	3	1	11	16	9	12	9	20	18	15	22	18	20	5	8	6
Capital and Coast	2	2	3	4	11	9	13	6	8	25	18	22	13	7	16	23	7	5	6	6
Counties Manukau	1	2	0	2	1	0	3	1	6	3	6	1	2	3	0	1	0	0	0	0
Hawke's Bay	0	0	1	3	1	1	3	0	4	4	8	2	3	8	7	3	1	2	3	2
Hutt Valley	4	0	0	0	0	1	1	1	0	0	0	0	0	1	0	0	0	0	0	0
Lakes	0	0	0	1	0	0	0	0	0	0	3	0	0	0	0	0	0	0	0	0
MidCentral	0	0	0	0	0	0	0	1	1	1	0	0	1	1	1	0	1	1	0	0
Nelson Marlborough	0	3	2	0	1	0	4	3	2	2	4	9	3	4	7	3	2	0	1	1
Northland	2	1	4	2	0	0	1	8	4	4	2	2	0	0	1	0	0	2	0	0
South Canterbury	1	3	1	2	0	0	2	1	1	1	0	5	10	8	8	16	23	6	6	9
Southern	1	2	0	5	4	5	1	2	4	4	7	11	4	10	11	4	6	5	6	4
Tairāwhiti	0	0	1	2	4	2	2	7	3	2	2	4	2	2	2	2	1	0	0	0
Taranaki	1	0	0	1	0	0	1	3	1	2	1	1	0	0	2	1	0	1	2	0
Waikato	0	5	3	3	4	6	4	8	4	13	9	6	5	3	7	4	1	7	2	0
Wairarapa	0	0	0	0	0	0	0	0	5	2	1	1	0	0	0	0	0	0	0	0
Waitemata	1	6	2	8	22	16	24	44	41	37	30	44	50	52	37	39	24	40	36	26
West Coast	1	1	0	0	0	2	0	1	3	2	0	0	0	1	2	0	0	0	1	1
Whanganui	0	0	1	4	0	1	2	0	5	5	5	0	6	4	1	1	1	0	0	0
<b>New Zealand</b>	<b>31</b>	<b>44</b>	<b>43</b>	<b>58</b>	<b>83</b>	<b>70</b>	<b>118</b>	<b>179</b>	<b>166</b>	<b>213</b>	<b>187</b>	<b>194</b>	<b>204</b>	<b>194</b>	<b>189</b>	<b>182</b>	<b>162</b>	<b>130</b>	<b>111</b>	<b>106</b>

**Table 6. Influenza-like illness rate by DHB by week 18–37, 2017**

DHB	Rate per 100 000																			
	18	19	20	21	22	23	24	25	26	27	28	29	30	31	32	33	34	35	36	37
Auckland	26.4	24.7	26.4	31.7	52.8	44.0	79.2	125.0	102.1	146.2	137.4	109.2	144.4	121.5	102.1	105.7	112.7	82.8	66.9	86.3
Bay of Plenty	0.0	0.0	0.0	0.0	13.5	6.7	6.7	40.4	47.1	74.1	26.9	26.9	33.7	40.4	47.1	47.1	74.1	60.6	13.5	13.5
Canterbury	2.8	7.1	14.2	4.3	4.3	1.4	15.6	22.8	12.8	17.1	12.8	28.4	25.6	21.3	31.3	25.6	28.4	7.1	11.4	8.5
Capital and Coast	7.8	7.8	11.7	15.6	42.8	35.0	50.6	23.4	31.2	97.3	70.1	85.7	50.6	27.3	62.3	89.6	27.3	19.5	23.4	23.4
Counties Manukau	3.0	6.0	0.0	6.0	3.0	0.0	9.0	3.0	18.0	9.0	18.0	3.0	6.0	9.0	0.0	3.0	0.0	0.0	0.0	0.0
Hawke's Bay	0.0	0.0	5.2	15.7	5.2	5.2	15.7	0.0	20.9	20.9	41.7	10.4	15.7	41.7	36.5	15.7	5.2	10.4	15.7	10.4
Hutt Valley	15.2	0.0	0.0	0.0	0.0	3.8	3.8	3.8	0.0	0.0	0.0	0.0	0.0	3.8	0.0	0.0	0.0	0.0	0.0	0.0
Lakes*	0.0	0.0	0.0	22.3	0.0	0.0	0.0	0.0	0.0	0.0	66.8	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0
MidCentral*	0.0	0.0	0.0	0.0	0.0	0.0	0.0	35.5	35.5	35.5	0.0	0.0	35.5	35.5	35.5	0.0	35.5	35.5	0.0	0.0
Nelson Marlborough	0.0	29.2	19.5	0.0	9.7	0.0	39.0	29.2	19.5	19.5	39.0	87.7	29.2	39.0	68.2	29.2	19.5	0.0	9.7	9.7
Northland	15.7	7.9	31.5	15.7	0.0	0.0	7.9	62.9	31.5	31.5	15.7	15.7	0.0	0.0	7.9	0.0	0.0	15.7	0.0	0.0
South Canterbury	9.3	27.8	9.3	18.6	0.0	0.0	18.6	9.3	9.3	9.3	0.0	46.4	92.8	74.2	74.2	148.5	213.4	55.7	55.7	83.5
Southern	2.2	4.4	0.0	10.9	8.7	10.9	2.2	4.4	8.7	8.7	15.3	24.0	8.7	21.8	24.0	8.7	13.1	10.9	13.1	8.7
Tairāwhiti	0.0	0.0	14.4	28.9	57.7	28.9	28.9	101.0	43.3	28.9	28.9	57.7	28.9	28.9	28.9	28.9	14.4	0.0	0.0	0.0
Taranaki	4.4	0.0	0.0	4.4	0.0	0.0	4.4	13.1	4.4	8.7	4.4	4.4	0.0	0.0	8.7	4.4	0.0	4.4	8.7	0.0
Waikato*	0.0	32.5	19.5	19.5	26.0	39.0	26.0	52.1	26.0	84.6	58.6	39.0	32.5	19.5	45.5	26.0	6.5	45.5	13.0	0.0
Wairarapa	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	200.0	80.0	40.0	40.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0
Waitematā*	7.9	47.5	15.8	63.3	174.0	126.6	189.8	348.0	324.3	292.7	237.3	348.0	395.5	411.3	292.7	308.5	189.8	316.4	284.7	205.6
West Coast	6.3	6.3	0.0	0.0	0.0	12.7	0.0	6.3	19.0	12.7	0.0	0.0	0.0	6.3	12.7	0.0	0.0	0.0	6.3	6.3
Whanganui*	0.0	0.0	60.2	241.0	0.0	60.2	120.5	0.0	301.2	301.2	301.2	0.0	361.4	241.0	60.2	60.2	60.2	0.0	0.0	0.0
<b>New Zealand</b>	<b>7.5</b>	<b>10.7</b>	<b>10.5</b>	<b>14.1</b>	<b>20.2</b>	<b>17.0</b>	<b>28.7</b>	<b>43.5</b>	<b>40.4</b>	<b>51.8</b>	<b>45.5</b>	<b>47.2</b>	<b>49.6</b>	<b>47.2</b>	<b>46.0</b>	<b>44.3</b>	<b>39.4</b>	<b>31.6</b>	<b>27.0</b>	<b>25.8</b>

\*Results that have some uncertainty, with less than 5% of the DHB population covered (see Notes on Interpretation).



Recent global experience with pandemic influenza A(H1N1)pdm09 highlights the importance of monitoring severe and mild respiratory disease to support pandemic preparedness as well as seasonal influenza prevention and control. Two active, prospective, population-based surveillance systems were used to monitor influenza and other respiratory pathogens: 1) among those registered patients seeking consultations with influenza-like illness (ILI) at sentinel general practices nation-wide; 2) among those hospitalized patients with severe acute respiratory illness (SARI) in Auckland and Counties Manukau District Health Boards (ADHB and CMDHB).

The aims of ILI and SARI surveillance are: 1) to measure the burden of severe and moderate disease caused by influenza and other respiratory pathogens; 2) to monitor trends in severe and moderate disease caused by influenza and other respiratory pathogens; 3) to identify high risk groups that should be prioritized for prevention and treatment; 4) to monitor antigenic, genetic and antiviral characteristics of influenza viruses associated with severe and mild disease. 5) to provide a study base to estimate the effectiveness of influenza vaccine.

### ACKNOWLEDGEMENT

We acknowledge the support of the New Zealand Ministry of Health. SARI surveillance was established and funded by the US CDC, and continues to operate through funding from the New Zealand Ministry of Health.

### DESCRIPTION OF ILI ACTIVITY THRESHOLDS

The values for the different intensity levels for 2017 are listed in the table below. This is based on New Zealand’s consultation rates from 2000–2015 (excluding the pandemic year, 2009) and WHO’s interim guidance severity assessment

Below seasonal level (baseline, per 100,000)	Seasonal level (per 100,000)			Above seasonal level (per 100,000)
	low	moderate	high	
<35.1	35.1-82.5	82.5-168.9	168.9-231.8	>231.8

- The baseline threshold indicates the level of influenza activity that signals the start and end of the annual influenza season and it is based on the Moving Epidemic Method (MEM) (*Vega et al. Influenza and other respiratory viruses 2013;7(4):546-558*).
- Seasonal levels (low, moderate and high) are estimated as the upper limits of the 40%, 90% and 97.5% one-sided confidence intervals of the geometric mean of 30 highest epidemic weekly rates using the MEM method. As many other countries use this method, it allows the NZ data to be interpreted not just at the country level but also comparable with other countries.
- The average seasonal curve indicates the usual seasonal activity that may occur during a typical year using the method described in “*Global epidemiological surveillance standards for influenza*” ([http://www.who.int/influenza/resources/documents/WHO\\_Epidemiological\\_Influenza\\_Surveillance\\_Standards\\_2014.pdf](http://www.who.int/influenza/resources/documents/WHO_Epidemiological_Influenza_Surveillance_Standards_2014.pdf)).

## NOTES ON INTERPRETATION

- SARI case definition: “An acute respiratory illness with a history of fever or measured fever of  $\geq 38^{\circ}\text{C}$ , AND cough, AND onset within the past 10 days, AND requiring inpatient hospitalisation (defined as a patient who is admitted under a medical team and to a hospital ward or assessment unit)”. A non-SARI case is a hospitalised respiratory patient who does not meet the SARI case definition.
- ILI case definition: “An acute respiratory illness with a history of fever or measured fever of  $\geq 38^{\circ}\text{C}$ , AND cough, AND onset within the past 10 days, AND requiring GP consultation”.
- ILI sentinel general practices: a total of 74 sentinel general practices have agreed to participate in community ILI surveillance. These practices have ~400 000 registered patients, covering roughly 9% of the NZ population.
- SARI sentinel hospitals serving a population of 906 000 people: Auckland City Hospital and the associated Starship Children’s Hospital (ADHB), and Middlemore Hospital and the associated Kidz First Children’s Hospital (CMDHB).
- The real-time PCR assay for influenza virus uses CDC’s protocol ([http://www.accessdata.fda.gov/cdrh\\_docs/pdf8/k080570.pdf](http://www.accessdata.fda.gov/cdrh_docs/pdf8/k080570.pdf));
- The real-time PCR assay for non-influenza respiratory viruses (respiratory syncytial virus, parainfluenza virus types 1-3, human metapneumovirus, rhinovirus and adenovirus) uses CDC’s protocol. Note: The rhinovirus PCR detects mostly rhinovirus with slight cross-reactivity against enterovirus.
- The surveillance week is Monday to Sunday inclusive, and data are extracted on the subsequent Tuesday. Results from previous weeks will be revised as data are updated (laboratory test results in particular may be delayed).
- ILI consultation rates for any particular DHB should be treated with caution. If the surveillance system has a small number of participating General Practices in the DHB, or the GP enrolled patient population is small, the calculated ILI rates are subject to greater fluctuation.

This weekly report is compiled by ESR. For more information please contact:

Tim Wood: T:+64 4 529 0611; E: [Tim.Wood@esr.cri.nz](mailto:Tim.Wood@esr.cri.nz)

Liza Lopez: T:+64 4 914 0647; E: [Liza.Lopez@esr.cri.nz](mailto:Liza.Lopez@esr.cri.nz)

Sue Huang: T:+64 4 529 0606; E: [Sue.Huang@esr.cri.nz](mailto:Sue.Huang@esr.cri.nz)